



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CD79A Recombinant Monoclonal Antibody [BLR234K]

Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_001774.1
Catalog No.	A700-234-T	Uniprot ID	P11912
Lot No.	A700-234-T-1	GeneID	973

APPLICATIONS WB, IP, IHC, ICC, Flow Cyt

SPECIES REACTIVITY Human, Mouse

AMOUNT 10 µl (5+ tests)

CONCENTRATION 500 µg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR234K

ORIGIN USA

PRODUCTION PROCEDURES Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 176 and the C-terminus (residue 226) of human B-cell antigen receptor complex-associated protein alpha chain (aka CD79A) using the numbering given in entry NP_001774.1 (Gene ID 973).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000

Immunoprecipitation 12 µl/1 mg lysate

Immunohistochemistry 1:50 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.

Immunocytochemistry 1:50 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.

Flow Cytometry Fixed in 4% formaldehyde and permeabilized with 90% methanol. 0.5 µl per 1×10^6 cells.


IHC HUMAN CONTROLS Appendix, Tonsil, Daudi Cells, Raji Cells, Ramos Cells

IHC MOUSE CONTROLS Spleen, CH27 Cells, WEHI-231 Cells


ADDITIONAL INFO <https://www.fortislifesciences.com/p/A700-234-T>

Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: September 29, 2022




Detection of human CD79A by western blot. *Samples:* Whole cell lysate (50 µg) from Daudi, HeLa, Raji, HEK293T, and Ramos cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).




Detection of mouse CD79A by western blot. *Samples:* Whole cell lysate (50 µg) from WEHI-231 (25 µg), NIH 3T3, CH27, TCMK-1, and A20 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).



Detection of human CD79A by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from Ramos cells prepared using NETN lysis buffer. *Antibodies:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234 lot 1) used for IP at 20 µl/mg lysate. CD79A was also immunoprecipitated by a second antibody against a different epitope of CD79A (A700-232). For blotting immunoprecipitated CD79A, A700-234 was used at 1:1000. Chemiluminescence with an exposure time of 1 second.



Detection of human CD79A by immunohistochemistry. *Sample:* FFPE section of human tonsil. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234-1). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).




Detection of mouse CD79A by immunohistochemistry.
Sample: FFPE section of mouse spleen. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234-1). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of human CD79A by immunocytochemistry.
Sample: FFPE section of Daudi cells. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234-1). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of mouse CD79A by immunocytochemistry. *Sample:* FFPE section of WEHI-231 cells.
Antibody: Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234-1). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of human CD79A (shaded) in Daudi cells by flow cytometry. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR234K] (A700-234) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120-101D5).