

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



#### **TIGIT Recombinant Monoclonal Antibody [BLR235K]**





Rabbit Recombinant Monoclonal

Purified RefSeq ID NP\_001139797.1

Catalog No. A700-235 Uniprot ID P86176
Lot No. A700-235-1 GeneID 100043314

APPLICATIONS WB

SPECIES REACTIVITY Mouse

**AMOUNT** 100 μl (10 blots)

CONCENTRATION 500 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

**ISOTYPE** IgG

CLONE # BLR235K
ORIGIN USA

JKIGIN 03A

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a recombinant protein representing the extracellular domain of mouse T-cell immunoreceptor with Ig and ITIM domains (TIGIT, residues 26–143, NP\_001139797.1).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000


ADDITIONAL INFO https://www.fortislife.com/p/A700-235

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: December 1, 2022



Detection of mouse TIGIT by western blot. Samples: Whole cell lysate (50 μg) from EL4, NIH 3T3, TCMK-1, ND7/23, and WEHI-231 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-TIGIT recombinant monoclonal antibody [BLR235K] (A700-235 lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 10 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).