



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



CD80 Recombinant Monoclonal Antibody [BLR237K]

Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_001346827.1
Catalog No.	A700-237	Uniprot ID	Q00609
Lot No.	A700-237-1	GeneID	12519


APPLICATIONS	WB, ICC, Flow Cyt
SPECIES REACTIVITY	Mouse
AMOUNT	100 µl (50+ tests)
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Borate Buffered Saline (BBS) pH 8.2 with 0.09% Sodium Azide, BSA Free
ISOTYPE	IgG
CLONE #	BLR237K
ORIGIN	USA
PRODUCTION PROCEDURES	Recombinant antibody was purified from cell culture supernatant. Immunogen was a recombinant protein representing the extracellular domain of mouse CD80 (residues 37–245, NP_001346827.1).
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections. Flow Cytometry Live or cells fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl per 1 × 10 ⁶ cells.
IHC MOUSE CONTROLS	C2C12 Cells, CH27 Cells
ADDITIONAL INFO	https://www.fortislife.com/p/A700-237 Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: December 19, 2022



Detection of mouse CD80 by western blot. *Samples:* Whole cell lysate (50 μ g) from C2C12, NIH 3T3, CH27, TCMK-1, and CT26 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-CD80 recombinant monoclonal antibody [BLR237K] (A700-237 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 3 minutes. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).

Detection of mouse CD80 by immunocytochemistry. *Sample:* FFPE section of C2C12 cells. *Antibody:* Rabbit anti-mCD80 recombinant monoclonal antibody [BLR237K] (A700-237). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of mouse CD80 (shaded) in CH27 cells by flow cytometry. *Antibody:* Rabbit anti-CD80 recombinant monoclonal [BLR237K] (A700-237) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120-101D5).