

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Troponin I Recombinant Monoclonal Antibody [BLR280L]



Rabbit Recombinant Monoclonal

Purified RefSeq ID NP_000354.4

Catalog No. A700-280-T Uniprot ID P19429 Lot No. 1 GeneID 7137

APPLICATIONS WB, IHC

SPECIES REACTIVITY Human, Mouse

AMOUNT 10 μl (5+ tests)

CONCENTRATION 250 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR280L
ORIGIN USA

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 0 and 50 of human

Troponin I (cardiac muscle) using the numbering given in entry NP_000354.4 (Gene ID 7137).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

IHC HUMAN CONTROLS Cardiac Muscle
IHC MOUSE CONTROLS Cardiac Muscle


ADDITIONAL INFO https://www.fortislife.com/p/A700-280-T


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD


Date: September 28, 2023






Detection of human Troponin I by western blot. Samples: Whole cell lysate (50 μg) from Jurkat, heart, and HEK293T cells prepared using NETN lysis buffer. Antibody: Rabbit anti-Troponin I recombinant monoclonal antibody [BLR280L] (A700–280 lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120–101P). Detection: Chemiluminescence with an exposure time of 3 seconds. Lower Panel: Rabbit anti-GAPDH antibody (A300–639A).

Detection of mouse Troponin I by western blot. Samples: Whole cell lysate (50 μg) from NIH 3T3, heart, and CH27 cells prepared using NETN lysis buffer. Antibody: Rabbit anti–Troponin I recombinant monoclonal antibody [BLR280L] (A700–280 lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti–rabbit IgG (A120–101P). Detection: Chemiluminescence with an exposure time of 3 seconds. Lower Panel: Rabbit anti–GAPDH antibody (A300–639A).



Detection of human Troponin I by immunohistochemistry. *Sample:* FFPE section of human cardiac muscle. *Antibody:* Rabbit anti-Troponin I recombinant monoclonal antibody [BLR280L) (A700–280). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120–501P).



Detection of mouse Troponin I by immunohistochemistry. Sample: FFPE section of mouse cardiac muscle. Antibody: Rabbit anti-Troponin I recombinant monoclonal antibody [BLR280L) (A700–280). Secondary: HRP-conjugated goat anti-rabbit IgG (A120–501P).