

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PVRL1/NECTIN1 Recombinant Monoclonal Antibody [BLR284L]



Rabbit Recombinant Monoclonal

Purified RefSeg ID NP_002846.3

Catalog No. A700-284 Uniprot ID Q15223 Lot No. 1 GeneID 5818

APPLICATIONS WB

SPECIES REACTIVITY Human, Mouse
AMOUNT 100 µl (10 blots)

CONCENTRATION 500 μg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR284L
ORIGIN USA

OKIGIN 03A

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 200 and 250 of human

Nectin-1 using the numbering given in entry NP_002846.3 (Gene ID 5818).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000


ADDITIONAL INFO https://www.fortislife.com/p/A700-284


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer. PhD

Date: September 28, 2023





Detection of human PVRL1/NECTIN1 by western blot. Samples: Whole cell lysate (10 μg) from MCF-10A, SNB-75, HaCaT, MCF-7, and RT-4 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-PVRL1/NECTIN1 recombinant monoclonal antibody [BLR284L] (A700-284 lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Detection: Chemiluminescence with an exposure time of 10 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal

antibody [BLR057F] (A700-057).

at lot 1) u anti-ra onds. Chemil nal Lower I

Detection of mouse PVRL1/NECTIN1 by western blot. Samples: Whole cell lysate (50 μg) from ND7/23, C6, RAW 264.7, J774A1, and AML12 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-PVRL1/NECTIN1 recombinant monoclonal antibody [BLR284L] (A700–284 lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120–101P). Detection: Chemiluminescence with an exposure time of 75 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700–057).