



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# IFN-gamma Recombinant Monoclonal Antibody [BLR300L]


Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_000610.2
Catalog No.	A700-300-T	Uniprot ID	P01579
Lot No.	1	GeneID	3458

---


<b>APPLICATIONS</b>	WB, IP, IHC, ICC
<b>SPECIES REACTIVITY</b>	Human
<b>AMOUNT</b>	10 µl (5+ tests)
<b>CONCENTRATION</b>	1000 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 - 8°C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Borate Buffered Saline (BBS) pH 8.2 with 0.09% Sodium Azide, BSA-Free
<b>ISOTYPE</b>	IgG
<b>CLONE #</b>	BLR300L
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Recombinant antibody was purified from cell culture supernatant.  Immunogen was a recombinant protein representing human Interferon gamma (residues 23-266, NP_000610.2).
<b>APPLICATIONS</b>	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.  Western Blot                      1:1,000 Immunoprecipitation            6 µl/1 mg lysate Immunohistochemistry        1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. Immunocytochemistry        1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.
<b>IHC HUMAN CONTROLS</b>	Appendix, Metastatic Lymph Node, Tonsil, HDLM-2 Cells
<b>ADDITIONAL INFO</b>	<a href="https://www.fortislife.com/p/A700-300-T">https://www.fortislife.com/p/A700-300-T</a> Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: September 28, 2023




**Detection of human IFN-gamma by western blot.**

*Samples:* Whole cell lysate (50 µg) from NK-92 cells untreated (-) and treated (+) with 80nM TPA, 3µM Ionomycin, 300ng/mL Brefeldin A. *Antibody:* Rabbit anti-IFN-gamma recombinant monoclonal antibody [BLR300L] (A700-300 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). *Detection:* Chemiluminescence with an exposure time of 3 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).




**Detection of human IFN-gamma by immunohistochemistry.**

*Sample:* FFPE section of tonsil. *Antibody:* Rabbit anti-IFN-gamma recombinant monoclonal antibody [BLR300L] (A700-300). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



**Detection of human IFN-gamma by western blot of immunoprecipitates.**

*Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from NK-92 cells treated with 80nM TPA, 3µM Ionomycin, and 300ng/mL Brefeldin A. *Antibodies:* Rabbit anti-IFN-gamma recombinant monoclonal antibody [BLR300L] (A700-300 lot 1) used for IP at 6 µl/mg lysate. IFN-gamma was also immunoprecipitated by a second antibody against a different epitope of IFN-gamma (Antibody 2). For blotting immunoprecipitated IFN-gamma, A700-300 was used at 1:1000. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



**Detection of human IFN-gamma by immunocytochemistry.**

*Sample:* FFPE section of NK-92 cells treated with TPA, Ionomycin, Brefeldin A (left) and mock treated NK-92 cells (right). *Antibody:* Rabbit anti-IFN-gamma recombinant monoclonal antibody [BLR300L] (A700-300). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).