



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Mouse IgG Heavy and Light Chain Cross-Adsorbed Antibody

Rabbit Polyclonal Conjugate DyLight® 488

Antigen Affinity Purified

Catalog No. A90-317D2

Lot No. A90-317D2-12

---

<b>APPLICATIONS</b>	WB, IHC, ICC, F, IF										
<b>SPECIES REACTIVITY</b>	Mouse. Minimum reactivity to human and rat										
<b>AMOUNT</b>	1 ml										
<b>CONCENTRATION</b>	0.5 mg/ml										
<b>STORAGE/SHELF LIFE</b>	2 – 8°C / 1 year from date of receipt										
<b>PHYSICAL STATE</b>	Liquid										
<b>BUFFER</b>	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide										
<b>FLUOROPHORE/PROTEIN</b>	7.9										
<b>ISOTYPE</b>	IgG										
<b>ORIGIN</b>	USA										
<b>PRODUCTION PROCEDURES</b>	<p>Antiserum was cross adsorbed using human and rat immunosorbents to remove cross reactive antibodies. The antibody to mouse IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to human and rat IgG was detected. This antibody may cross react with IgG from other species.</p>										
<b>APPLICATIONS</b>	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.</td></tr><tr><td>Immunohistochemistry</td><td>1:40 – 1:400</td></tr><tr><td>Immunocytochemistry</td><td>1:40 – 1:500</td></tr><tr><td>Flow Cytometry</td><td>1:50 – 1:200</td></tr><tr><td>Immunofluorescence</td><td>1:40 – 1:500</td></tr></table>	Western Blot	1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.	Immunohistochemistry	1:40 – 1:400	Immunocytochemistry	1:40 – 1:500	Flow Cytometry	1:50 – 1:200	Immunofluorescence	1:40 – 1:500
Western Blot	1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.										
Immunohistochemistry	1:40 – 1:400										
Immunocytochemistry	1:40 – 1:500										
Flow Cytometry	1:50 – 1:200										
Immunofluorescence	1:40 – 1:500										
<b>APPLICATION NOTES</b>	<p>Not all listed applications have been specifically tested by our laboratory.</p> <p>DyLight® 488 is excited at 493 (in PBS) and emits at 518 (in PBS).</p> <p>DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.</p>										
<b>ADDITIONAL INFO</b>	<p><a href="https://www.bethyl.com/product/A90-317D2">https://www.bethyl.com/product/A90-317D2</a></p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>										

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.


Michael Spencer, PhD

Date: June 10, 2022


Phone: 800.338.9579 • Fax: 866.597.6105 • Web: [www.bethyl.com](http://www.bethyl.com)  
Orders: [orders@fortislife.com](mailto:orders@fortislife.com) • Support: [technical@fortislife.com](mailto:technical@fortislife.com)

1





**Detection of GAPDH and hnRNP in HeLa Whole Cell Lysate.** *Primary Antibodies:* cocktail of goat anti-GAPDH A303-878A (A303-878A-1) and mouse anti-hnRNP A500-011A (A500-011A-1) at 1 µg/ml each. *Secondary Antibodies:* cocktail of Dylight® 680-conjugated rabbit anti-goat A50-200D6 (A50-200D6-1) (red) and Dylight® 488-conjugated rabbit anti-mouse A90-317D2 (A90-317D2-3) (blue) at 0.5 µg/ml each. *Acquisition:* Syngene G:Box, 39 seconds (red) and 60 seconds (blue).



**Detection of human p53 by immunofluorescence.** *Sample:* FFPE section of human breast carcinoma. *Primary Antibody:* mouse anti-p53 (clone DO-1) used at a dilution of 1:100. *Secondary Antibody:* Green-fluorescent Rabbit anti-mouse IgG-heavy and light chain cross-adsorbed Antibody DyLight® 488 Conjugated (A90-317D2 Lot 6) used at a dilution of 1:100 (5µg/ml). *Counterstain:* DAPI (blue)