



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Catalogue

 nordicmubio.com/products/p16-ink4a-dcs-50/P16002-L

p16 INK4a (DCS-50)

Catalog number: **P16002-L**

Clone	DCS-50
Isotype	IgG1
Product Type	Monoclonal Antibody Primary Antibodies
Units	1ml
Host	Mouse
Species Reactivity	Human.
Application	Immunohistochemistry (frozen & paraffin) Immunoprecipitation Western Blotting

Background

The antibody is suitable to detect p16 in different tissues. Staining results may be cytoplasmatic or nuclear. Nuclear staining should be more specific. Human p16 INK4a (syn. CDKN2; MTS-1) protein is an inhibitor of Cyclin-Dependent-Kinases 4 and 6 (cdk4/cdk6). CDK4 is an important enzyme for the progression of the cell cycle during the G1-Phase. By its inhibitory action p16 is an important regulator in the cell cycle. The gene encoding for p16 is deleted in many tumour cell lines. In adenocarcinoma of the cervix uteri usually p16 levels are increased. Endocervical p16-overexpression is mostly associated with "high risk" HPV-Typing. In other tumour types as squamous epithelial carcinoma in cervix uteri and head and neck carcinoma p16 expression usually is reduced.

Source

Monoclonal antibody DCS-50 is produced after immunisation of Balb/c mice with the full length recombinant p16 INK4a protein of human origin.

Immunogen: Recombinant human p16 INK4a molecule

Product

Protein A affinity purified antibody lyophilized from PBS pH7.4 with BSA and Na-Azide 0.09%.

Purification Method: Protein A affinity purification.

Secondary Reagents: We recommend the use of BIOLOGO's Universal Staining System DAB (Art. No. DA005) or AEC (Art.-No. AE005).

Concentration: Approximately 50 µg/ml after reconstitution in 1 ml distilled water

Specificity

p16 INK4a

Species Reactivity: Human, other species not tested

Applications

IHC (C,P), IP

Incubation Time: 60 min at RT

Working Concentration: 1:10

Pre-Treatment: Pre-treatment with Unmasking Fluid C (Citrate Buffer, Art. No. DE000) or Unmasking Fluid G (Art. No. DE007) at 96-100 degree of Celsius is recommended for paraffin sections.

Positive Control: Colon carcinoma, HPV-pos. cervical carcinoma

Storage

2-8°C

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. It may contain hazardous ingredients. Please refer to the Safety Data Sheets (SDS) for additional information and proper handling procedures. Dispose product remainders according to local regulations. This datasheet is as accurate as reasonably achievable, but Exalpha Biologicals accepts no liability for any inaccuracies or omissions in this information.

References

1. Koh J., Enders G.H., Dynlacht B.D., and Harlow E. (1995) Tumour-derived p16 alleles encoding proteins defective in cell cycle inhibition. *Nature* 375; 506-510. 2. Reed A.L., Califano J., Cairns P., Westra W.H., Jones R.M., et al. (1996) High frequency of p16 (CDKN2/MTS-1/INK4A) inactivation of head and neck squamous cell carcinoma. *Cancer Res.* 56(16); 3630-3633. 3. Geradts J., Hruban R.H., Schütte M., Kern S.E., and Maynard R. (2000) Immunohistochemical p16INK4a analysis of archival tumors with deletion, hypermethylation, or mutation of the CDKN2/MTS1 gene. A comparison of four

commercial antibodies. Appl. Immun 4. Negri G., Egarter-Vigl E., Kasal A., Romano F., Haitel A., Mian C. (2003) p16INK4a is a useful marker for the diagnosis of adenocarcinoma of the cervix uteri and its precursors: an immunohistochemical study with immunocytochemical correlations. Am J. S 5. Kotaro R. Shibata, Tomoki Aoyama, Yasuko Shima et al. (2007) Expression of the p16INK4A Gene Is Associated Closely with Senescence of Human Mesenchymal Stem Cells and Is Potentially Silenced by DNA Methylation During In Vitro Expansion Stem Cells vol. 9; 2371–2382.

Protein Reference(s)

Database Name: UniProt

Accession Number: P42771 (CD2A1_HUMAN)

Species Accession: Human