



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)


[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## NAD<sup>+</sup>, free acid

**Catalog No:** 16110  
**Lot No:** XXXXX  
**Cas No:** 53-84-9  
**Formula:** C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>  
**MW:** 663.43  
**Supplied as:** solid  
**Stability:** store at -20°C



### Background

Nicotinamide adenine dinucleotide, abbreviated NAD<sup>+</sup>, is a coenzyme found in all living cells. The compound is a dinucleotide, since it consists of two nucleotides joined through their phosphate groups. One nucleotide contains an adenine base and the other nicotinamide. In metabolism, NAD<sup>+</sup> is involved in redox reactions, carrying electrons from one reaction to another. The coenzyme is, therefore, found in two forms in cells: NAD<sup>+</sup> is an oxidizing agent – it accepts electrons from other molecules and becomes reduced. This reaction forms NADH, which can then be used as a reducing agent to donate electrons. These electron transfer reactions are the main function of NAD<sup>+</sup>. However, it is also used in other cellular processes, the most notable one being a substrate of enzymes that add or remove chemical groups from proteins, in posttranslational modifications. Because of the importance of these functions, the enzymes involved in NAD<sup>+</sup> metabolism are targets for drug discovery.

### Tests

#### Appearance:

#### Appearance of solution:

#### Purity (HPLC):

#### β-NAD (Enzym, 340 nm):

#### Water content (K.F.):

#### Absorbtion

**A<sub>250</sub>/A<sub>260</sub>**

**A<sub>280</sub>/A<sub>260</sub>**

### Specifications

almost white to faint yellowish

lyophilized powder

clear, colourless to faint yellowish

≥95%

≥94.5%

≤3.5%

0.81 – 0.85

0.20 – 0.24

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**