



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Activin A, His Tag, human recombinant (rHuActivin-A-His)

**Catalog No:** 40707  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** Activin Beta-A chain, Erythroid differentiation protein, EDF, FRP, Activin-A, Inhibin-B, Inhibin-Beta A chain

### Background

Inhibins are dimeric peptide hormones produced by female ovarian granulosa cells and male Sertoli cells as well as a variety of other tissues. Inhibins have two isoforms, A and B, with the same alpha subunit but different beta subunits. Inhibin A is a dimer of alpha and beta A subunits, inhibin B is a dimer of alpha and beta B subunits. Inhibins are thought to inhibit the production of follicle-stimulating hormone (FSH) by the pituitary gland. In addition, Inhibins are also thought to play a role in the control of gametogenesis, and embryonic and fetal development.

### Description

Activin-A human recombinant also called Inhibin-Beta A chain produced in *E. coli* is a single monomeric, non-glycosylated, polypeptide chain containing 116 amino acids (311-426) and having an N-terminal hexa-histidine tag, having a total molecular weight of 17.47 kDa. Activin-A is purified by standard chromatographic techniques.

### Physical Appearance

Sterile filtered clear solution.

### Formulation

Activin-A is supplied in 20 mM Tris-HCl, 1 mM EDTA and 50% glycerol.

### Stability

Store Inhibin-Beta Alpha chain at 4°C if entire vial will be used within 2-4 weeks. Store frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

### Purity

Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**