

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







Interleukin-1 beta, mouse recombinant (rmIL-1-beta)

Catalog No: 50441 Lot No: XXXXX Source: *E. coli*

Synonyms: Catabolin, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator

(LEM), Mononuclear Cell Factor (MCF), IL1F2, IL-1 beta

Background

Interleukin-1b is produced by activated macrophages, IL-1B stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1B proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

Description

Interleukin-1 beta mouse recombinant produced in *E. coli* is a non-glycosylated, polypeptide chain containing 153 amino acids and having a molecular mass of 17500 Dalton. IL-1b is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Mouse IL-1b was lyophilized from sterile filtered aqueous solution 10 mM sodium phosphate, pH 7.5.

Solubility

It is recommended to reconstitute the lyophilized Interleukin-1b in sterile 18 M Ω -cm H $_2$ O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Interleukin-1b, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1b should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95.0% as determined by SDS-PAGE.

Amino Acid Sequence

MVPIRQLHYR LRDEQQKSLV LSDPYELKAL HLNGQNINQQ VIFSMSFVQG EPSNDKIPVA LGLKGKNLYL SCVMKDGTPT LQLESVDPKQ YPKKKMEKRF VFNKIEVKSK VEFESAEFPN WYISTSQAEH KPVFLGNNSG QDIIDFTMES VSS

Activity

The activity as determined by the dose-dependant stimulation of mouse D10S cells is 0.9 - 1.3 pg/ml, corresponding to a specific activity of 1.1×10^6 IU/mg.





Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.