



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Interleukin-7, mouse recombinant (rmlL-7)

**Catalog No:** 50447  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** Lymphopoietin 1 (LP-1), pre-B cell factor, IL-7

### Background

Interleukin-7 is a potent lymphoid cell growth factor produced primarily by stromal cells. IL-7 has been shown to support the proliferation and differentiation of pre B- and early T cells as well as displaying a biological effect on cells of NK and myeloid lineages.

### Description

Interleukin-7 mouse recombinant produced in *E. coli* is a single, non-glycosylated polypeptide chain containing 129 amino acids and having a molecular mass of 14.9 kDa. IL-7 is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

Filtered (0.2 µm) and lyophilized from a concentrated (1 mg/ml) solution in 1×PBS, pH 7.4 and 2% trehalose.

### Solubility

It is recommended to reconstitute the lyophilized Interleukin-7 in sterile 18 MΩ-cm H<sub>2</sub>O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized Interleukin-7, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-7 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 96.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

### Amino Acid Sequence

ECHIKDKEGK AYESVLMISI DELDKMTGTD SNCPNNEPNF FRKHVCD DTK EAAFLNRAAR KLKQFLKMNI SEEFNVHLLT  
VSQGTQTLVN CTSKEEKNVK EQKKN DACFL KRLLEIKTC WNKILK GSI

### Activity

The ED<sub>50</sub> as determined by the dose-dependant stimulation of murine 2E8 cells is <0.2 ng/ml, corresponding to a specific activity of ≥5,000,000 units/mg.50447

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**