



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Angiotensin-like Protein-4, His Tag, human recombinant (rHuANGPTL-4-His)

Catalog No: 87332
Lot No: XXXXX
Source: *E. coli*
Synonyms: ANGPTL4, NL2, ARP4, FIAF, PGAR, HFARP, pp1158, ANGPTL2, Fasting- Induced Adipose Factor, Hepatic Fibrinogen/Angiotensin-Related Protein, PPARG Angiotensin-Related Protein

Background

FIAF (fasting-induced adipose factor) a.k.a ANGPTL4 or PGAR or HFARP is an adipocytokine up-regulated by fasting, by peroxisome proliferator-activated receptor agonists, and by hypoxia. ANGPTL4 is found in human and mouse blood plasma both as a native protein and in a truncated form. In human white adipose tissue and SGBS adipocytes, only the native form of ANGPTL4 could be detected, whereas in mice the differentiation of mouse 3T3-L1 adipocytes is associated with the production of truncated ANGPTL4. However, the truncated ANGPTL4 is produced by human liver. In human blood plasma FIAF is mainly presented in a truncated form (FIAF-S2), whose levels fenofibrate treatment increases (as shown by experimental data). There is an inter individual variation in ANGPTL4 levels of both the truncated and the native form, however those levels were not influenced by prolonged semistarvation and are not associated with body mass index.

Description

ANGPTL4 human recombinant is manufactured with N-terminal fusion of His Tag. The Angiotensin-like protein 4 His - tagged fusion protein is 25 kDa protein containing 204 amino acid residues of Angiotensin-like protein 4 and 16 additional amino acid residues - His Tag. The amino acid sequence of the ANGPTL4 human recombinant is 100% homologous to the 26-229 amino acid sequence of the Angiotensin-like protein-4 precursor without signal sequence. ANGPTL4 is purified by one-step procedure using affinity Ni-NTA chromatography.

Physical Appearance

Filtered white lyophilized (freeze-dried) powder.

Formulation

Filtered and lyophilized from 0.5 mg/ml in 0.05 M acetate buffer pH 4.

Solubility

Add 0.2 ml of 0.1 M acetate buffer pH4 and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10 µg/ml.

Stability

Store lyophilized Angiotensin-like Protein 4 at -20°C. Aliquot the ANGPTL4 after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted ANGPTL4 can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Purity

Greater than 95% as determined by SDS-PAGE.



Amino Acid Sequence

MRGSHHHHHH GMASHMGPVQ SKSPRFASWD EMNVLAHGLL QLQQLREHA ERTRSQLSAL ERRLSACGSA CQGTEGSTDL
PLAPESRVDP EVLHSLQTQL KAQNSRIQQL FHKVAQQQRH LEKQHLRIQH LQSQFGLLDH KHLDEHAVAKP ARRKRLPEMA
QPVDPAHNVS RLHRLPRDCQ ELFQVGERQS GLFEIQPQGS PPFLVNCKMT SDGGWTVIQR

Applications

WB*ELISA

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.