



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Hepatocyte Growth Factor, human recombinant (rHuHGF)

**Catalog No:** 94893  
**Lot No:** XXXXX  
**Source:** Insect cells  
**Synonyms:** Scatter Factor (SF), Hepatopoietin (HPTA), HGF, HGFB, F-TCF

### Background

Hepatocyte Growth Factor (HGF) is a multifunctional growth factor which regulates both cell growth and cell motility. It exerts a strong mitogenic effect on hepatocytes and primary epithelial cells. HGF synergizes with Interleukin-3 and GM-CSF to stimulate colony formation of hematopoietic progenitor cells in vitro and may, therefore, also modulate hematopoiesis.

### Description

HGF human recombinant produced in Baculovirus is a heterodimer, non-glycosylated, polypeptide chain containing 692 amino acids and having a total molecular mass of 78 kDa. HGF is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

The sterile protein powder (1 mg/ml) is lyophilized from a solution containing 50 mM acetic acid.

### Solubility

It is recommended to reconstitute the lyophilized HGF in 50 mM acetic acid to a concentration not less than 100 µg/ml. Further dilutions should be made using buffer containing protein.

### Stability

Lyophilized Hepatocyte Growth Factor, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution HGF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 95.0% as determined by SDS-PAGE.

### Amino Acid Sequence

QRKRRNTIHE FKKSAKTTLI KIDPALKIKT KKVNTADQCA NRCTRNGKLP FTCKAFVFDK ARKQCLWFPP NSMSSGVKKE  
FGHEFDLYEN KDYIRNCIIG KGRSYKGTVS ITKSGIKCQP WSSMIPHEHS YRGKDLQENY CRNPRGEEGG PWCFTSNPEV  
RYEVCDDIPQC SEVEMTCNGE SYRGLMDHTE SGKICQRWDH QTPHRHKFLP ERYPDKGFDD NYCRNPDGQP RPWCYTLDPH  
TRWEYCAIKT CADNTMNDTD VPLETTECIQ GQEGYRGTV NTIWNIGIPCQ RWDSQYPHEH DMTPENFKCK DLRENYCRNP  
DGSESPWCFT TDPNIRVGYC SQIPNCDMSH GQDCYRGNGK NYMGNLSQTR SGLTCSMWDK NMEDLHRHIF WEPDASKLNE  
NYCRNPDDDA HGPWCYTGNP LIPWDYCPIS RCEGDTTPTI VNLDPVISC AKTKQLRVVN GIPTRTNIGW MVSLRYYRNKH  
ICGGLIKES WVLTAQCFCP SRDLKDYEAW LGIHDVHGRG DEKCKQVLNV SQLVYGPEGS DLVLMKLARP AVLDDFVSTI  
DLPNYGCTIP EKTSCSVYGW GYTGLINYDG LLRVAHLYIM GNEKCSQHRH GKVTLNESEI CAGAEEKIGSG PCEGDYGGPL  
VCEQHKMRMV LGVIVPGRGC AIPNRPGIFV RVAYYAKWIH KIILTYKVPQ S



**Activity**

The activity was assayed for scattering activity in the MDCK cell assay. The ED50 for this effect is typically at 1.0 - 5.0 ng/ml (200,000 - 1,000,000 IU/mg).

**Usage**

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**