



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Hepatocyte Growth Factor, human recombinant (rHuHGF-CHO)

**Catalog No:** 94897  
**Lot No:** XXXXX  
**Source:** CHO cells  
**Synonyms:** Scatter Factor (SF), Hepatopoietin (HPTA), HGF

### Background

Hepatocyte Growth Factor (HGF) is a multifunctional growth factor which regulates both cell growth and cell motility. It exerts a strong mitogenic effect on hepatocytes and primary epithelial cells. HGF synergizes with Interleukin-3 and GM-CSF to stimulate colony formation of hematopoietic progenitor cells in vitro and may, therefore, also modulate hematopoiesis.

### Description

HGF human recombinant produced in CHO is a heterodimer, non-glycosylated, polypeptide chain consisting of an a-chain of 463 amino acids and b-chain of 234 having a total molecular mass of 75 kDa. HGF is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

The protein was lyophilized from a concentrated (1.4 mg/ml) solution containing phosphate buffered saline with 0.02% Tween 80, pH 7.4.

### Solubility

It is recommended to reconstitute the lyophilized HGF in sterile 18 M $\Omega$ -cm H<sub>2</sub>O not less than 100  $\mu$ g/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized Hepatocyte Growth Factor, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution HGF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 97.0% as determined by SDS-PAGE.

### Amino Acid Sequence

Agrees with the sequence of native human HGF.

### Activity

The ED<sub>50</sub>, calculated by the dose-dependant proliferation of monkey 4 MBr-5 indicator cells was found to be 20 - 40 ng/ml corresponding to a specific activity of 25,000 - 50,000 IU/mg.

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**