



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Granulocyte Colony Stimulating Factor, mouse recombinant (rmG-CSF)

**Catalog No:** 94967  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** CSF3, MGI-1G, GM-CSF beta, Pluripoietin, G-CSF, GCSF

### Background

Granulocyte Colony Stimulating Factor is a growth factor and/or cytokine produced by the endothelium, macrophages and a number of other immune cells. GCSF stimulates the bone marrow to produce granulocytes and also to stimulate the survival, proliferation, differentiation and function of neutrophil granulocyte progenitor cells and mature neutrophils.

### Description

Granulocyte Colony Stimulating Factor mouse recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 178 amino acids and having a molecular mass of 18.9 kDa. G-CSF is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

GCSF was lyophilized from 10 mM Sodium citrate, pH 4.0 and 150 mM NaCl.

### Solubility

It is recommended to reconstitute the lyophilized GCSF in sterile 18 M $\Omega$ -cm H<sub>2</sub>O not less than 100  $\mu$ g/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized Granulocyte Colony Stimulating Factor, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution GCSF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 98.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

### Amino Acid Sequence

VPLVTVSAL PPSLPLPRSF LLKSLEQVRK IQASGSVLE QLCATYKLCH PEELVLLGHS LGIPKASLSG CSSQALQQTQ  
CLSQHLHSGLC LYQGLLQALS GISPALAPTL DLLQLDVANF ATTIWQQMEN LGVAPTQPT QSAMPAFTSA FQRRAGGVLA  
ISYLQGFLET ARLALHHLA

### Activity

The ED50 calculated by the dose-dependant proliferation of murine NFS-60 indicator cells (measured by <sup>3</sup>H-thymidine uptake) is <0.05 ng/ml, corresponding to a specific activity of 2 x 10<sup>7</sup> IU/mg.



**Usage**

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.