



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Macrophage Migration Inhibitory Factor, His Tag, human recombinant (rHuMIF-His-C)

Catalog No: 95053
Lot No: XXXXX
Source: *E. coli*
Synonyms: Phenylpyruvate tautomerase, Glycosylation-inhibiting factor, GIF, MMIF, MIF

Background

The cytokine Macrophage migration inhibitory factor (MIF) has been identified to be secreted by the pituitary gland and the monocyte/macrophage and to play an important role in endotoxic shock. MIF has the unique property of being released from macrophages and T cells in response to physiological concentrations of glucocorticoids. The secretion of MIF is tightly regulated and decreases at high, anti-inflammatory steroid concentration.

Description

MIF human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 123 amino acids and having a molecular mass of 13.5 kDa. MIF is fused to a His tag at C-terminus and purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered lyophilized powder.

Formulation

Human MIF was lyophilized from a 1 mg/ml solution containing PBS pH 7.4.

Solubility

It is recommended to reconstitute the lyophilized MIF in sterile 18 M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized MIF, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution MIF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

MPMFIVNTNV PRASVPDGF^L SELTQQLAQA TGKPPQYIAV HVVPDQLMAF GGSSEPALCS LHSIGKIGGA QNRSYSKLLC
GLLAERLRIS PDRVYINYD MNAANVGWNN STFAL^EHHHH HH

Activity

Measured by its ability to bind rhCD74 in a functional ELISA.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.