

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







Interferon alpha2a, His Tag, human recombinant, tobacco (rHuIFN-alpha2a-His)

Catalog No: 97075 Lot No: XXXXX

Source: Nicotiana benthamiana

Synonyms: Leukocyte interferon, B cell interferon, Type I interferon, IFNA2, IFN-a 2a

Background

IFN-alpha is produced by macrophages and has antiviral activities. Interferon stimulates the production of two enzymes: protein kinase and an oligoadenylate synthetase.

Description

Interferon alpha 2a human recombinant produced in plant is a single, glycosylated, polypeptide chain containing 165 amino acids and having a molecular mass of 19 kDa. Interferon-a 2a contains a 6xHis Tag at C-terminus. IFN-A 2a is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized, containing 50% SDS.

Solubility

It is recommended to reconstitute the lyophilized IFN-a 2a in sterile 18 M Ω -cm H $_2$ O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Interferon alpha 2a, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN-alpha 2a should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 99.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

The sequence of the first twelve N-terminal amino acids was determined and was found to be Cys-Asp-Leu-Pro-Gln-Thr-His-Ser-Leu-Gly-Ser-Arg.

Activity

The specific activity was determined by a quantitative gene report bioassay using human Type I interferon-sensitive cells. Compared with bacterial derived recombinant IFN-a 2a, assayed in the same conditions, plant derived IFN-alpha 2a showed 6.1×10^8 units/mg in reference to a viral resistance assay using bovine kidney MDBK cells.





Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.