



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Lymphotactin (XCL1), His Tag, human recombinant (rHuXCL1-His)

**Catalog No:** 97223  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** XCL1, Cytokine SCM-1, ATAC, Lymphotaxin, SCM-1-alpha, Small inducible cytokine C1, XC chemokine ligand 1, LTN, LPTN, SCM1, SCM-1, SCYC1, SCM-1a

### Background

XCL1 is a small cytokine belonging to the XC chemokine family that is also known as lymphotactin. It is located in high levels in spleen, thymus, intestine and peripheral blood leukocytes, and at lower levels in lung, prostate gland and ovary. Cellular sources for XCL1 include activated thymic and peripheral blood CD8+ T cells. This chemokine attracts T cells. In humans, XCL1 is closely related to another chemokine called XCL2, whose gene is found at the same locus on chromosome 1. XCL1 induces its chemotactic function by binding to a chemokine receptor called XCR1.

### Description

XCL1 human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 114 amino acids (22-114) and having a molecular mass of 12.5 kDa. XCL1 is fused to a 20 amino acid His Tag at N-terminus and is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered colorless solution.

### Formulation

Recombinant XCL1 contains 20 mM Tris buffer pH 8, 200 mM NaCl, 2 mM DTT and 30% glycerol.

### Stability

XCL1, although stable at 10°C for 3 weeks, should be stored desiccated below -18°C. Please prevent freeze-thaw cycles.

### Purity

Greater than 95.0% as determined by SDS-PAGE.

### Amino Acid Sequence

MGSSHHHHHH SSSLVPRGSH MVGSEVSDKR TCVSLTTQRL PVSRIKTYTI TEGSLRAVIF ITKRGLKVCA DPQATWVRDV  
VRSMDRKSNT RNNMIQTKPT GTQQSTNTAV TLTG

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**