



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Interleukin-17A/F Heterodimer, human recombinant (rHuIL-17A/F)

**Catalog No:** 97225  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** IL17A/F, IL17 A/F, IL-17A/F, IL-17 A/F, IL17AF, IL-17 AF, Interleukin-17 A/F, Interleukin-17 AF

### Background

Human IL-17A/F is a 40 kDa glycoprotein which is secreted as a disulfide-linked heterodimer. IL-17A/F consists of two proteins of the IL-17 family, IL-17A and IL17F. Proteins of the 6 homodimeric IL17 family show a cysteine knot motif that contains two disulfide-bonds. Human IL17A is produced as a 155 a.a precursor that includes a 23 amino acids signal sequence and a 132 amino acid chain that includes an N-linked glycosylation site. Human IL17F is produced as a 153 amino acid precursor with a 20 amino acid signal sequence and a 133 amino acid region. Similar to IL17A, IL17F also has an N-linked glycosylation site. Both proteins (IL17A & IL17F) share 50% amino acid sequence identity. Human IL17A & IL17F show approximately 60% homology in their amino acid sequence to mouse IL-17A and IL-17F. Interleukin-17A/F and IL17A, IL17F homodimers are manufactured by activated CD4+ T cells, called Th17. IL-23 causes Th17 lymphocytes to manufacture IL-17A/F. IL17RA and IL17RC form a heterodimer for the binding of IL17A and IL17F. IL-17A/F binds IL-17RA. Interleukin-17A/F induces chemokine production and airway neutrophilia with intermediate potency between IL17A (most potent) and IL17F (least potent).

### Description

Interleukin-17A/F human recombinant produced in *E. coli* is a heterodimeric, non-glycosylated polypeptide chain containing 1 monomeric subunit of each IL-17A and IL-17F. The active dimer contains 271 amino acids and having a total molecular mass of 30.7 kDa. IL-17A/F is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

Lyophilized from a concentrated (1 mg/ml) solution containing no additives.

### Solubility

It is recommended to reconstitute the lyophilized Interleukin-17A/F in sterile water not more than 1 mg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized IL-17A/F, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Human IL-17A/F should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 98.0% as determined by SDS-PAGE.



### Amino Acid Sequence

MIVKAGITIP RNPGPCNSED KNFPRTVMVN LNIHNRNTNT NPKRSSDYNN RSTSPWNLHR NEDPERYPSV IWEAKCRHLG  
CINADGNVDY HMNSVPIQQE ILVLRREPPH CPNSFRLEKI LVSVGCTCVT PIVHHVAMRK IPKVGHTFFQ KPESCPPVPG  
GSMKLDIGII NENQRVSMSR NIESRSTSPW NYTVTWDPNR YPSEVVQAQC RNLGCINAQG KEDISMNSVP IQQETLVVRR  
KHQGCVSFQ LEKVLVTVGC TCVTPVIHHV Q

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**