



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Fibroblast Growth Factor 12 (FGF12), His-Tag, human recombinant (rHuFGF12-His)

Catalog No: 97346
Lot No: XXXXX
Source: *E. coli*
Synonyms: FGF-12, FGF12, FGF12B, FHF1, Fibroblast growth factor 12, Fibroblast growth factor homologous factor 1, FHF-1, Myocyte-activating factor

Background

FGF12 is part of the Fibroblast Growth Factor (FGF) family which has a vast mitogenic and cell survival functions, and play a role in a range of biological activities, among them are embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. FGF-12 doesn't obtain the N-terminal signal sequence present in the majority of the FGF family members, but it contains clusters of basic residues that act as a nuclear localization signal. When transfected into mammalian cells, FGF12 accumulated in the nucleus, but was not secreted. FGF12 is involved in nervous system development and function. FGF12 binds to IB2 (islet brain-2), a cellular kinase scaffold, and voltage gated sodium channels and is also involved in intracellular signaling and ion exchange.

Description

FGF-12 human recombinant protein is a single, non-glycosylated polypeptide chain produced in *E. coli*, having a molecular weight of 22.6 kDa and containing 201 amino acids (1-181). FGF12 is fused to a 20 amino acid His Tag at the N-terminus.

Formulation

The FGF-12 solution (1 mg/ml) contains 20 mM Tris pH 7.5, 1 mM DTT, 2 mM EDTA and 10% glycerol.

Stability

Store FGF12 at -20°C. Can be stored at 4°C for a limited period of time of 7 days.

Purity

Greater than 90% as determined by SDS-PAGE.

Amino Acid Sequence

MSSHHHHHHS SGLVPRGSHM ESKEPQLKGI VTRLFSQQGY FLQMHPDGTI DGTKDENS DY TLFNLI PVGL RVVAIQGVKA
SLYVAMNGEY LYSSDVFTPE CKFKESVFEN YYVIYSSTLY RQQESGRAWF LGLNKEGQIM KGNRVKKTTP SHFVPKPIEV
CMYREQLSHE IGEKQGRKRS SGTPTMNGGK VVNQDST

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.