



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Transforming Growth Factor-Beta 2, His Tag, human recombinant (rHuTGF-b2-His)

Catalog No: 97384
Lot No: XXXXX
Source: *Nicotiana benthamiana*
Synonyms: Transforming growth factor, beta 2, cetermin, Glioblastoma-derived T-cell suppressor factor, polyergin, G-TSF, TGF-beta2, TGF-beta-2, transforming growth factor beta-2, BSC-1 cell growth inhibitor, TGFB-2

Background

TGFB2 is a 27.08 kDa protein having two identical 118 amino acid peptide chains linked by a single disulfide bond. TGFB2 is part of a family of five related cytokines that have an extensive variation of normal and neoplastic cells, indicating the importance of these homo-dimer proteins as multi-functional regulators of cellular activity. The three mammalian isoforms of TGF- β (TGF β 1, TGF β 2 and TGF β 3) signal through the same receptor and stimulate similar biological responses. They are involved in physiological processes as embryogenesis, tissue remodelling and wound healing.

Description

TGF-b2 human recombinant produced in plants is a homodimeric polypeptide chain containing 2 x 118 amino acids and having a total molecular mass of 27.08 kDa. TGFB2 is fused to 6xHis Tag at N-terminus and purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a concentrated (1 mg/ml) solution containing 50 mM Tris-HCl pH 7.4.

Solubility

It is recommended to reconstitute the lyophilized TGFB2 in sterile 18 M Ω -cm H₂O not less than 1 μ g/40 μ l, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized TGFB2, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TGFB2 Human should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 97.0% as determined by SDS-PAGE.

Amino Acid Sequence

HHHHHHALDA AYCFRNVQDN CCLRPLYIDF KRDLGWKWIH EPKGYNANFC AGACPYLWSS DTQHRSVLSL YNTINPEASA
SPCCVSDLE PLTILYYIGK TPKIEQLSNM IVKSCKCS

Activity

The biological activity of TGFB2 is measured in culture by its ability to inhibit the mink lung epithelial (Mv1Lu) cells proliferation. ED50 <40 ng/ml, corresponding to a specific activity of 25,000 units/mg.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.