



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Beta Defensin-3, mouse recombinant (rmBD-3)

Catalog No: 97413
Lot No: XXXXX
Source: *E. coli*
Synonyms: Beta-defensin 3, BD-3, mBD-3, Defensin beta 3, Defb3, Bd3, MGC129397

Background

Defensins form a family of microbicidal and cytotoxic peptides made by neutrophils. Members of the defensin family are highly similar in protein sequence. This gene encodes defensin, beta 103A, which has broad spectrum antimicrobial activity and may play an important role in innate epithelial defense.

Description

Beta Defensin-3 mouse recombinant produced in *E. coli* is a single, non-glycosylated polypeptide chain containing 41 amino acids and having a molecular mass of 4.6 kDa. BD-3 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Mouse BD-3 was lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.

Solubility

It is recommended to reconstitute the lyophilized Beta Defensin-3 in sterile 18 MΩ-cm H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Beta Defensin-3, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution BD-3 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Purity

Greater than 98.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

KKINNPVSCL RKGGRWCNRC IGNTRQIGSC GVPFLKCKCR K

Activity

Measured by its antimicrobial activity against *E. coli*. The ED₅₀ for this effect is typically 4 - 20 µg/ml, corresponding to a specific activity of 50,000 - 250,000 units/mg.

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.