



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)





## Interleukin-1 beta, HEK, human recombinant (rHuIL-1 beta-HEK)

**Catalog No:** 97484

**Lot No:** XXXXX

**Source:** HEK293

**Synonyms:** Catabolin, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF), IL1F2, IL-1 beta

### Background

Interleukin-1b is produced by activated macrophages, IL-1B stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1B proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

### Description

Interleukin-1 beta human recombinant produced in HEK cells is a glycosylated monomer, having a molecular weight range of 18-25 kDa due to glycosylation. IL-1 beta is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

### Formulation

The IL-1 beta was lyophilized from 1 mg/ml in 1xPBS.

### Solubility

It is recommended to reconstitute the lyophilized Interleukin-1b in sterile PBS containing 0.1% endotoxin-free recombinant HSA.

### Stability

Lyophilized IL-1 beta, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-1 beta should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Purity

Greater than 95% as observed by SDS-PAGE.

### Activity

The specific activity was determined by the dose-dependent stimulation of the proliferation of mouse D10S cells and is typically 0.02 - 0.08 ng/ml.

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**