



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Interleukin-23, His Tag, HEK, human recombinant (rHuIL-23-His-HEK)

**Catalog No:** 97503  
**Lot No:** XXXXX  
**Source:** HEK293  
**Synonyms:** Interleukin 23 alpha subunit p19, Interleukin-12 subunit beta p40, SGRF, IL23P19, IL-23-A, interleukin-six, G-CSF related factor, JKA3 induced upon T-cell activation, interleukin 12B (natural killer cell stimulatory factor 2 cytotoxic lymphocyte maturation factor 2 p40), NK cell stimulatory factor chain 2, Cytotoxic lymphocyte maturation factor 40 kDa subunit, CLMF2, CLMF p40

### Background

IL23 is composed of a subunit of the heterodimeric cytokine IL23 and the p40 subunit of interleukin 12 (IL12B). Interleukin-23 (IL-23) belongs to the IL-12 family and is produced by antigen presenting cells. IL-23 using IL12RB1 and IL-23R (specific for IL-23) can activate STAT and NF- $\kappa$ B pathways and stimulate the production of interferon-gamma. IL-23 is known to take a vital part in the inflammation process and is associated with auto immune diseases. However, unlike IL12, which acts primarily on naive CD4(+) T cells, IL23 preferentially acts on memory CD4(+) T cells.

### Description

Interleukin-23 human recombinant produced in HEK 293 cells is a heterodimer (IL23A co-transfected with IL12B), containing one polypeptide chain of IL23 protein (19.77 kDa) fused with a C-terminal His Tag and one polypeptide chain of IL12b-Flag (35.69 kDa). IL-23 is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered clear solution.

### Formulation

IL-23 His Tag protein is supplied in 50 mM Tris pH 7.5, 130 mM NaCl and 10% glycerol.

### Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

### Purity

Greater than 90.0% as determined by analysis by SDS-PAGE.

### Amino Acid Sequence

IL23A-His :

RAVPGGSSPA WTQCQQLSQK LCTLAWSAHP LVGHMDLREE GDEETTNDVP HIQCGDGDGP QGLRDNSQFC LQRIHQGLIF  
YEKLLGSDIF TGEPSLLPDS PVGQLHASLL GLSQLLQPEG HHWETQQIPS LSPSQPWQRL LLRFKILRSL QAFVAVAAARV  
FAHGAATLSP HHHHHHHH

IL12B-Flag :

IWELKKDVYV VELDWYPDAP GEMVVLTCDT PEEDGITWTL DQSSEVLGSG KTLTIQVKEF GDAGQYTCHK GGEVLSHSL  
LLHKKEDGIW STDILKDQKE PKNKTFLRCE AKNYSGRFTC WWLTTISTDL TFSVKSSRGS SDPQGVTCGA ATLSAERVRG  
DNKEYEYSVE CQEDSACPAA EESLPIEVMV DAVHKLKYEN YTSSFFIRDI IKPDPKLNQ LKPLKNSRQV EVSWEYPDTW  
STPHSYFSLT FCVQVQGKSK REKKDRVFTD KTSATVICRK NASISVRAQD RYSSSSWSEW ASVPCSDYKD DDDK



**Usage**

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.