



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Macrophage Inflammatory protein-3 alpha (CCL20), His Tag, human recombinant (rHuMIP-3a-His)

Catalog No: 97539
Lot No: XXXXX
Source: *E. coli*
Synonyms: S Small inducible cytokine A20 precursor, CCL20, Macrophage inflammatory protein 3 alpha, MIP-3-alpha, Liver and activation- regulated chemokine, CC chemokine LARC, Beta chemokine exodus-1, CKb4, LARC, ST38, MIP3A, MIP-3a, SCYA20

Background

CCL-20 is a chemotactic factor that draws lymphocytes & neutrophils, rather than monocytes. MIP-3 alpha inhibits proliferation of myeloid progenitors in colony formation assays. MIP3A plays a role in the formation and function of the mucosal lymphoid tissues by attracting lymphocytes and dendritic cells towards epithelial cells. C-terminal processed forms have been shown to be equally chemotactically active for leukocytes. CCL-20 holds antibacterial activity *E. coli* atcc 25922 and *s.aureus* atcc 29213. CCL-20 gene transcription is activated by *H. pylori*, which activates NF-kappaB through intracellular signal pathway which involves I kappaB kinase and NF-kappaB-inducing kinase. MIP-3 alpha is involved in chemokine-mediated lymphocyte trafficking during gastric inflammation in *Helicobacter* infection. CCL-20 expression is involved in the recruitment of CD45RO-positive T cell subsets into the intestinal lamina propria. MIP-3A is in charge of the advancement of pulpal inflammation through the recruitment of C-C motif Receptor 6-expressing lymphocytes. Vaginal epithelial cells respond to factors present in semen by secreting MIP-3 alpha, which increases langerhans cells recruitment during HIV transmission.

Description

Macrophage Inflammatory protein-3 alpha human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 91 amino acids (27-96) and having a molecular mass of 10.3 kDa. MIP 3a is fused with a 21 amino acid His Tag at N-terminus and purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered colorless solution.

Formulation

The MIP 3a solution (0.25 mg/ml) contains phosphate buffered saline pH 7.4 and 10% glycerol.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Purity

Greater than 85.0% as determined by SDS-PAGE.

Amino Acid Sequence

MGSSHHHHHHH SSSLVPRGSH MASNFDCCLG YTDRLHPKF IVGFTRQLAN EGCINAIIF HTKKKLSVCA NPKQTWVKYI
VRLLSKKVKN M

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.