



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Vascular Endothelial Growth Factor D, His Tag, HEK, human recombinant (rHuVEGF-D-His -HEK)

Catalog No: 97582
Lot No: XXXXX
Source: HEK293
Synonyms: c-fos induced growth factor (vascular endothelial growth factor D), FIGF, VEGFD

Background

VEGF-D belongs to the VEGF/PDGF family of proteins. VEGF-D promotes lymphangiogenesis, endothelial cell growth, and regulates vascular permeability. In addition, VEGF-D has an important part in the creation of the venous and lymphatic vascular systems and in the growth and maintenance of differentiated lymphatic endothelium. Mature VEGF-D forms a noncovalently linked homodimer, and binds to and activates both VEGFR-2 (flk1) and VEGFR-3 (flt4).

Description

VEGFD human recombinant produced in HEK-293 cells is a secreted protein (amino acids Phe93-Ser201) fused to a polyhistidine tag at the C-terminus.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Recombinant VEGF-D was lyophilized after extensive dialysis against PBS.

Solubility

It is recommended to reconstitute the VEGF D in sterile 18 M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized VEGF-D, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution VEGF-D should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95.0% as determined by SDS-PAGE.

Activity

The ED50 of 3-4 ng/ml is measured by its ability to stimulate the proliferation of human microvascular endothelial cells (HMVECs).

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.