

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







Platelet Derived Growth Factor-D, His tag, human recombinant (rHuPDGFD-His)

Catalog No: 97588
Lot No: XXXXX
Source: E. coli

Synonyms: Platelet Derived Growth Factor D, Spinal Cord-Derived Growth Factor B, Iris-Expressed Growth Factor,

SCDGF-B, IEGF, PDGF-D, MSTP036

Background

Platelet-derived growth factor D (PDGFD) belongs to the platelet-derived growth factor family. PDGFD gene product only forms homodimers and, thus, does not dimerize with the other 3 family members. PDGFD has an imperative role in wound healing. PDGFD induces macrophage recruitment, increased interstitial pressure, and blood vessel maturation during angiogenesis. PDGFD initiates events which lead to a mesangial proliferative glomerulonephritis, including influx of monocytes and macrophages and production of extracellular matrix. The 4 members of the PDGF family are mitogenic factors for cells of mesenchymal origin and are distinguished by a core motif of eight cysteines, 7 of which are found in this factor. PDGFD differs from alpha and beta members of this family by having an odd N-terminal domain, the CUB domain.

Description

PDGFD human recombinant produced in *E. coli* is a single polypeptide chain containing 146 amino acids (250-370) and having a molecular mass of 16.6 kDa. PDGFD is fused to a 25 amino acid His Tag at N-terminus and purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered colorless solution.

Formulation

The PDGFD solution (1 mg/1 ml) contains 20 mM Tris-HCl buffer (pH 8.0), 0.4 M urea and 10% glycerol.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Purity

Greater than 90% as determined by SDS-PAGE.

Amino Acid Sequence

MGSSHHHHHH SSGLVPRGSH MGSHMSYHDR KSKVDLDRLN DDAKRYSCTP RNYSVNIREE LKLANVVFFP RCLLVQRCGG NCGCGTVNWR SCTCNSGKTV KKYHEVLQFE PGHIKRRGRA KTMALVDIQL DHHERCDCIC SSRPPR

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.