



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Interleukin-13, rhesus macaque recombinant (rrmIL-13)

Catalog No: 97595
Lot No: XXXXX
Source: *E. coli*
Synonyms: NC30, ALRH, BHR1, P600, IL-13, MGC116786, MGC116788, MGC116789

Background

IL13 is an immunoregulatory cytokine produced primarily by activated Th2 cells. IL-13 is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4.

Description

Interleukin-13 Rhesus Macaque recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 114 amino acids and having a molecular mass of 12.6 kDa. IL 13 Rhesus Macaque is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4 containing 5% trehalose.

Solubility

It is recommended to reconstitute the lyophilized IL-13 in sterile 18 MΩ-cm H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized IL-13, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-13 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Purity


Greater than 97.0% as determined by SDS-PAGE and HPLC analyses.

Amino Acid Sequence

SPSPVPRSTA LKELIEELVN ITQNQKAPLC NGSMVWSINL TAGVYCAALE SLINVSGCSA IEKTQRMING FCPHKVSAGQ
FSSLRVRDTK IEVAQFVKDL LVHLKKLFRF GRFN

Activity

The ED50 as determined by the dose-dependent proliferation of TF-1 cells was <1.0 ng/ml, corresponding to a specific activity of >1 × 1,000,000 units/mg.



Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.