



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Angiotensin-like Protein 2, human recombinant (rhuARP2)

**Catalog No:** 99928  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** Angiotensin-related protein 2, Angiotensin-like protein 2, ANGPTL2, ARP2, HARP

### Background

Angiotensins belong to the vascular endothelial growth factor family and are the only known growth factors largely specific for vascular endothelium. Angiotensins-1, -2 and -4 partake in the formation of blood vessels. ANGPTL2 displays angiogenic effects. Angiotensin-like protein 2 (ANGPTL2) is an anti-diabetic factor. It induces sprouting in endothelial cells through autocrine and paracrine action. ANGPTL2 increases insulin sensitivity in adipocytes. In addition, ANGPTL2 is a mediator of chronic adipose tissue inflammation. ANGPTL2 is widely expressed in the heart, small intestine, spleen and stomach. ANGPTL2 is also found in lower levels in the colon, ovary, adrenal gland, skeletal muscle and prostate.

### Description

ANGPTL2 Human Recombinant, produced in *E. coli*, is a single, non-glycosylated, polypeptide chain containing 481 amino acids including a 10 amino acid N-terminal His tag. The total molecular mass is 56 kDa (calculated).

### Physical Appearance

Filtered, white lyophilized (freeze-dried) powder.

### Formulation

ANGPTL2 was filtered (0.4 µm) and lyophilized from 0.5 mg/ml in 0.03 M acetate buffer (pH 4.0).

### Solubility

It is recommended to add 0.1 M acetate buffer (pH 4) to prepare a working stock solution of approximately 0.5 mg/ml and let the lyophilized pellet dissolve completely. For conversion into higher pH value, an intensive dilution by relevant buffer to a concentration of 10 g/ml is recommended. In higher concentrations the solubility of the ANGPTL2 is limited. ANGPTL2 is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

### Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time. It does not show any change after two weeks at 4°C.

### Amino Acid Sequence

MKHHHHHHAS QEDGFEGTEE GSPREFIYLN RYKRAGESQD KCTYTFIVPQ QRVTGAIQVN SKEPEVLLEN RVHKQELELL  
NNELLKQKRQ IETLQQLVEV DGGIVSEVKL LRKESRMNS RVTQLYMQLL HEIIRKRDNA LELSQLENRI LNQTADMLQL  
ASKYKDLEHK YQHLATLAHN QSEIIAQLLE HCQRVPSARP VPQPPPAAPP RVYQPPTYNR IINQISTNEI QSDQNLKVL P  
PPLPTMPTLT SLPSTDKPS GPWRDCLQAL EDGHDTSIY LVKPENTNRL MQVWCDQRHD PGGWTVIQRR LDGSVNFFRN  
WETYKQGFNG IDGEYWLGLE NIYWLTNQGN YKLLVTMEDW SGRKVFAEYA SFRLEPESEY YKLRLGRYHG NAGDSFTWHN  
GKQFTTLDRD HDVYTGNAH YQKGGWYNA CAHSNLNGVW YRGGHYRSRY QDGVYWAEFR GGSYSLKKVV MMIRPNPNTF H

### Usage

**This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.**