



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Interleukin 17B, human recombinant (rhIL20)

**Catalog No:** 99934  
**Lot No:** XXXXX  
**Source:** *E. coli*  
**Synonyms:** Interleukin-17B, IL-17B, Cytokine Zcyto7, Interleukin-20, Neuronal interleukin-17-related factor, IL20, NIRF, ZCYTO7

### Background

IL-17 family members are glycoproteins secreted as dimers, which induce local cytokine production and recruit granulocytes to sites of inflammation. The IL-17 family is comprised of at least six pro-inflammatory cytokines that share a conserved cysteine-knot structure but diverge at the N-terminus. IL-17 is induced by IL-15 and IL-23, mostly in activated CD4+ T cells distinct from Th1 or Th2 cells. IL-17B binds the IL-17B receptor, but not the IL-17 receptor. It is most homologous with IL-17D, which is expressed by resting CD4+ T cells and CD19+ B cells. Diseases associated with IL17B include spondyloarthritis and neuronitis, and among its related super-pathways are Mucin expression in CF via IL-6, IL-17 signaling pathways and STAT3 Pathway.

### Description

Interleukin-17B Human Recombinant, produced in *E. coli*, is a homodimeric, non-disulfide-linked polypeptide chain containing a total of 322 amino acids (2 chains of 161 aa). It has a molecular mass of 36.5 kDa. The IL-17B is purified by proprietary chromatographic techniques.

### Physical Appearance

Sterile filtered, white, lyophilized (freeze-dried) powder.

### Formulation

Interleukin-17B was lyophilized from a 0.2 µm filtered concentrated solution in PBS (pH 7.4), 0.05% Tween-20 and 3% trehalose.

### Solubility

It is recommended to reconstitute the lyophilized Interleukin 17B in sterile 18 MΩ-cm H<sub>2</sub>O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

### Stability

Lyophilized Interleukin 17B, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL17B should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Amino Acid Sequence

MQPRSPKSKR KGQGRPGPLA PGPHQVPLDL VSRMKPYARM EYERNIEEM VAQLRNSSEL AQRKCEVNLQ LWMSNKRSL  
PWGYSINHDP SRIPVDLPEA RCLCLGCVNP FTMQEDRSMV SVPVFSQVPV RRRLCPPPPR TGPCRQRAVM ETIavgctci F

### Activity

The ED<sub>50</sub>, as determined by inducing IL-8 secretion of human HepG2 cells, is less than 1.0 µg/ml, corresponding to a specific activity of > 1000 IU/mg.



**Usage**

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.