



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Data Sheet

TGF β R1, Fc fusion, Avi-Tag

Human, Recombinant, C-terminal Fc-Avi-tag
Catalog #: 100644
Lot #: 200220 **Conc.:** 0.45 mg/ml

Formulated in: 8 mM Phosphate, pH 7.4,
110 mM NaCl, 2.2 mM KCl, and 20%
glycerol.

Stability: At least 6 months at -80°C . *Avoid
freeze/thaw cycles.*

References:

1. Loeys, B.L., *et al.* *Nature Genetics*.
2005; **37**: 275-281.
2. Singh, K.K., *et al.*, *Human Mutation*.
2006; **27(8)**: 770-777.

Description: Human Transforming growth factor beta receptor I (TGF β R1), also known as activin A receptor type II-like kinase, AAT5 and LDS1, GenBank Accession No. NM_004612.4, a.a 34-126, fused at the C-terminus of the Fc portion of human IgG1, with C-terminal Avi-tagTM, expressed in a HEK293 cell expression system. MW=39 kDa. This protein runs at a higher molecular weight due to glycosylation.

Quality Assurance


4-20% SDS-PAGE Coomassie staining

Lane 1:
4 μ g TGF β R1

Lane 2:
Protein Marker

MW: 39 kDa + glycans

Purity: $\geq 90\%$



OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**
Or you can Email us at: info@bpsbioscience.com
Please visit our website at: www.bpsbioscience.com