



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Product Information


Description:	Recombinant anti-TFR (Transferrin Receptor), which recognizes TFR. The purified antibody was enzymatically biotinylated using Avi-tag™ technology. This affinity purified antibody has been tested for specific binding to purified human TFR1 (BPS Bioscience #101287) in an ELISA binding assay.
Species	Human
Concentration:	0.64 mg/ml
Expression System:	HEK293
Purity:	≥90%
Format:	Aqueous buffer solution.
Formulated In:	8 mM phosphate, pH 7.4, 110 mM NaCl, 2.2 mM KCl, and 20% glycerol
MW:	Heavy Chain: 50 kDa; Light Chain: 24 kDa + glycans
Glycosylation:	This antibody runs at a higher MW by SDS-PAGE due to glycosylation.
Label:	This antibody is enzymatically biotinylated using Avi-Tag™ technology. Biotinylation is confirmed to be ≥90%.
Stability:	At least 12 months at -80°C.
Storage:	-80°C
Instructions for Use:	Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before opening. Aliquot into small volumes and flash freeze for long term storage. Avoid multiple freeze/thaw cycles.
Assay Conditions:	An ELISA binding assay was performed with this antibody and its target protein, TFR1 (BPS Bioscience #101287). The target protein was coated onto an assay plate overnight at 2 µg/ml at 4°. The next day, the plate was washed on blocked. Then, serial dilutions of the antibody were added (0 to 200 nM) for 1 hour at room temperature. After washing, HRP-Streptavidin was incubated for 30 minutes at room temperature. Following another wash step, colorimetric HRP substrate was added and quenched with 1N HCL. Then we measured absorbance on a plate reader at 450 nm. EC50 = 0.61 nM
Applications:	Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.

Quality Control Data

4-20% SDS-PAGE Coomassie Staining




TFR1:Anti-TFR (Biotin) Activity



Quality Control Data

Biotin-Avidin Pulldown



1. Beads
2. Flow thru
3. Control
4. Standards

* Avidin from beads.