



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product Information

Description:	Recombinant human FGFR1a (IIIc) (fibroblast growth factor receptor 1 alpha IgIII domain c), encoding amino acids 22-376, corresponding to the extracellular domain. This construct has a C-terminal human IgG1 Fc domain followed by an Avi-Tag™. This protein was affinity purified.
Background:	FGFR1 (fibroblast growth factor receptor 1), also known as FLT2, is a member of the FGFR family, a family of four membrane-spanning tyrosine kinases (FGFR1-4) which serve as high-affinity receptors for 17 growth factors (FGF1-17). The FGF Receptor family plays an important role in multiple biological processes, including mesoderm induction and patterning, cell growth and migration, organ formation and bone growth. FGFR1 is alternatively spliced generating multiple splice variants that are differentially expressed during embryo development and in the adult body. It undergoes splicing on the IgIII domain, the domain involved in ligand specificity, creating two alternative forms: IIIb and IIIc. FGFR1a (IIIc) can bind to FGF1, 2, 4, 5, 6, 19, 20, and 21. FGFR proteins are tightly regulated, being usually ubiquitinated by CBL (Cbl proto-oncogene E3 ubiquitin protein ligase) after activation, with FGFR1 being more susceptible to undergo ubiquitination. FGFR1 has been linked to breast, ovarian, and other types of cancer, making the development of FGFR1-targeted inhibitors an active and promising field of research in cancer therapy.
Species:	Human
Construct:	FGFR1 (22-376-Fc(IgG1)-Avi)
Concentration:	2.09 mg/ml
Expression System:	HEK293
Purity:	≥90%
Format:	Aqueous buffer solution.
Formulated In:	8 mM phosphate, pH 7.4, 110 mM NaCl, 2.2 mM KCl, and 20% glycerol
MW:	69 kDa + glycans
Glycosylation:	This protein runs at a higher MW by SDS-PAGE due to glycosylation.
Genbank Accession:	NM_023110.3
Stability:	At least 6 months at -80°C.
Storage:	-80°C
Instructions for Use:	Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before opening. Aliquot into small volumes and flash freeze for long term storage. Avoid multiple freeze/thaw cycles.
Applications:	Useful for SDS-PAGE.

Quality Control Data

4-20% SDS-PAGE Coomassie Staining

