



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Product Information

|                              |  |
|------------------------------|--|
| <b>Description:</b>          | Complex of recombinant human full length CDK1 (cyclin dependent kinase 1) and CyclinE1. Both constructs contain an N-terminal GST-tag. The complex was affinity purified and is active.  |
| <b>Species:</b>              | Human  |
| <b>Construct:</b>            | CDK1 (GST-Full Length) / Cyclin E1 (GST-Full Length)   |
| <b>Concentration:</b>        | 0.10 mg/ml   |
| <b>Expression System:</b>    | Co-expressed in Sf9  |
| <b>Purity:</b>               | ≥90%   |
| <b>Format:</b>               | Aqueous buffer solution.   |
| <b>Formulated In:</b>        | 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM Glutathione, 0.1 mM EDTA, 0.25 mM DTT, and 25% glycerol   |
| <b>MW:</b>                   | CDK1: 58 kDa<br>Cyclin E1: 73 kDa  |
| <b>Genbank Accession:</b>    | CDK1: NM_001786<br>Cyclin E1: NM_001238  |
| <b>Stability:</b>            | At least 6 months at -80°C.  |
| <b>Storage:</b>              | -80°C  |
| <b>Instructions for Use:</b> | Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before opening. Aliquot into small volumes and flash freeze for long term storage. Avoid multiple freeze/thaw cycles.   |
| <b>Specific Activity:</b>    | 244 pmol/min/μg  |
| <b>Assay Conditions:</b>     | CDK1/CyclinE1 activity was measured by using Histone H1 peptide (GGPATP-KKAKKL-COOH), diluted in 20 mM Tris-HCl (pH 7.5) to a working concentration of 1 mg/ml, in the ADP Glo™ Kinase Assay kit (Promega #V9101). Reaction was initiated by mixing increasing amounts of CDK1/CyclinE1 with 25 μM ATP in 40 mM Tris-HCl, pH 7.4, 20 mM MgCl <sub>2</sub> , 0.1 mg/ml BSA prepared with 50 μM DTT, and substrate at a final concentration of 200 μg/ml.<br>After a 40-minute incubation at room temperature, the reaction was terminated by addition of ADP-Glo™ Reagent, followed by a subsequent 40-minute incubation at room temperature. Kinase Detection Reagent was added, and the reaction was incubated for another 30 minutes at ambient temperature. Detection of luminescence was measured using the Luminescence Module Protocol on GloMax®-Multi Microplate Multimode Reader. The Specific Activity was calculated as follows: (Corrected activity, RLU) / [(Specific activity from ADP in RLU/pmol) * (Reaction time in min) *(Enzyme amount in μg or mg)]. Corrected RLU was calculated by subtracting the blank value from all the values. The blank was determined from a “no enzyme” sample by replacing the enzyme solution with an equal volume of Kinase Dilution Buffer IX (1x). |
| <b>Applications:</b>         | Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.  |

## Quality Control Data

4-20% SDS-PAGE Coomassie Staining



Specific Activity

