



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product Information

Description:	Complex of recombinant human full length CDK2 (cyclin dependent kinase 2) and CyclinE2. Both constructs contain an N-terminal GST-tag. The complex was affinity purified and is active.
Species:	Human
Construct:	CDK2 (GST-Full Length) / Cyclin E2 (GST-Full Length)
Concentration:	0.10 mg/ml
Expression System:	Co-expressed in Sf9
Purity:	≥90%
Format:	Aqueous buffer solution.
Formulated In:	50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM Glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, and 25% glycerol
MW:	CDK2: 58 kDa Cyclin E2: 72 kDa
Genbank Accession:	CDK2: NM_001798 Cyclin E2: NM_057749
Stability:	At least 6 months at -80°C.
Storage:	-80°C
Instructions for Use:	Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before opening. Aliquot into small volumes and flash freeze for long term storage. Avoid multiple freeze/thaw cycles.
Specific Activity:	16 pmol/min/μg
Assay Conditions:	CDK2/CyclinE2 activity was measured by using Histone H1 peptide (GGGPATP-KKAKKL-COOH), diluted in 20 mM Tris-HCl (pH 7.5) to a working concentration of 1 mg/ml, in the ADP Glo™ Kinase Assay kit (Promega #V9101). Reaction was initiated by mixing increasing amounts of CDK2/CyclinE2 with 25 μM ATP in 40 mM Tris-HCl, pH 7.4, 20 mM MgCl ₂ , 0.1 mg/ml BSA prepared with 50 μM DTT, and substrate at a final concentration of 200 μg/ml. After a 40-minute incubation at room temperature, the reaction was terminated by addition of ADP-Glo™ Reagent, followed by a subsequent 40-minute incubation at room temperature. Kinase Detection Reagent was added, and the reaction was incubated for another 30 minutes at ambient temperature. Detection of luminescence was measured using the Luminescence Module Protocol on GloMax®-Multi Microplate Multimode Reader. The Specific Activity was calculated as follows: $\text{Specific Activity} = \frac{(\text{Corrected activity, RLU})}{[(\text{Specific activity from ADP in RLU/pmol}) * (\text{Reaction time in min}) * (\text{Enzyme amount in } \mu\text{g or mg})]}$ Corrected RLU was calculated by subtracting the blank value from all the values. The blank was determined from a “no enzyme” sample by replacing the enzyme solution with an equal volume of Kinase Dilution Buffer IX (1x).
Applications:	Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.

Quality Control Data

4-20% SDS-PAGE Coomassie Staining



Specific Activity

