



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet
BRDT (BD1+BD2) TR-FRET Assay Kit
Catalog # 32616
Size: 384 reactions

DESCRIPTION:

The BRDT (BD1+BD2) TR-FRET Assay Kit is designed to measure the inhibition of BRDT (BD1+BD2) binding to its substrate in a homogeneous 384 reaction format. This FRET-based assay requires no time-consuming washing steps, making it especially suitable for high throughput screening applications. The assay procedure is straightforward and simple; a sample containing terbium-labeled donor, dye-labeled acceptor, BRDT, ligand, and an inhibitor is incubated for 120 minutes. Then, the fluorescence intensity is measured using a fluorescence reader.

COMPONENTS:

| Catalog # | Component | Amount | Storage | |
|-----------|---|-----------|------------|-------------------------------------|
| 31061 | BRDT (BD1+BD2) | 10 µg | -80°C | (Avoid freeze/ thaw cycles!) |
| 33000 | BET Bromodomain Ligand | 50 µl | -80°C | |
| | Non-acetylated Ligand 1 | 15 µl | -80°C | |
| | Tb donor | 2 x 10 µl | -20°C | |
| | Dye-labeled acceptor | 2 x 10 µl | -20°C | |
| 33012 | 3x BRD TR-FRET Assay Buffer 1 | 4 ml | -20°C | |
| | White, Nonbinding, low volume, microtiter plate | 1 | Room temp. | |

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Fluorescent microplate reader capable of measuring Time Resolved Fluorescence Resonance Energy Transfer (TR-FRET)
Adjustable micropipettor and sterile tips

APPLICATIONS: Great for screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: At least 6 months from date of receipt when stored as directed.

REFERENCE(S): Filippakopoulos, P., *et al.*, *Cell* 2012; **149**:214.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**
Or you can Email us at: info@bpsbioscience.com
Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

Protocol for BRDT (BD1+BD2) assay

- 1) Dilute one part **3x BRD TR-FRET Assay Buffer 1** with 2 parts distilled water (3-fold dilution) to make **1x BRD TR-FRET Assay Buffer 1**. Make only a sufficient quantity needed for the assay; store remaining stock solution in aliquots at -20°C.
- 2) Dilute **Tb-labeled donor** and **Dye-labeled acceptor** 100-fold in **1x BRD TR-FRET Assay Buffer 1**. Make only sufficient quantities needed for the assay; store remaining stock solution in aliquots at -20°C.
- 3) Add 5 µl of diluted **Tb-labeled donor**, and 5 µl of diluted **Dye-labeled acceptor** to each well designated "Test Inhibitor", "Negative Control", and "Positive Control".
- 4) Add 2 µl of inhibitor solution to each well designated "Test Inhibitor". Add 2 µl of the same solution without inhibitor (inhibitor buffer) to the wells labeled "Negative Control", and "Positive Control".

| | Negative Control* | Positive Control | Test Inhibitor |
|---------------------------------|-------------------|------------------|----------------|
| Tb-labeled donor | 5 µl | 5 µl | 5 µl |
| Dye-labeled acceptor | 5 µl | 5 µl | 5 µl |
| Test Inhibitor | - | - | 2 µl |
| Inhibitor Buffer (no inhibitor) | 2 µl | 2µl | - |
| BET Bromodomain Ligand | | 5 µl | 5 µl |
| Non-acetylated Ligand 1 | 5 µl | - | - |
| BRDT (BD1 + BD2) 3 ng/µl | 3 µl | 3 µl | 3 µl |
| Total | 20 µl | 20 µl | 20 l |

***Non-acetylated Ligand 1** may be used as a substrate control in place of the negative control

- 5) Thaw **BET Bromodomain Ligand** and **Non-acetylated Ligand 1** on ice. Upon first thaw, briefly spin tube containing enzyme to recover the full contents of the tube. Aliquot each ligand into single-use aliquots. Store remaining undiluted ligand at -80°C immediately. *Note: each ligand is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots.*
- 6) Individually dilute **BET Bromodomain** 40-fold in **1x BRD TR-FRET Assay Buffer 1**. Add 5 µL of diluted **BET Bromodomain Ligand** to each well designated as "Positive Control" and "Test Inhibitor". Add 5 µL of **1x BRD TR-FRET Assay Buffer 1** to the wells labeled as "Negative Control". *Note: if using the **Non-acetylated Ligand 1**, dilute **Non-acetylated Ligand 1** 40-fold in **1x BRD TR-FRET Assay Buffer 1** and add 5 µl of diluted **Non-***
OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

acetylated Ligand 1 to the “Negative Control” well in place of the 5 μ l of **1x TR-FRET BRD Assay Buffer 1**.

- 7) Thaw **BRDT** bromodomain protein on ice. Upon first thaw, briefly spin tube containing protein to recover the full contents of the tube. Aliquot **BRDT** protein into single-use aliquots. Store remaining undiluted **BRDT** in aliquots at -80°C immediately. *Note: BRDT is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted protein.*
- 8) Dilute **BRDT** in **1x BRD TR-FRET Assay Buffer 1** to 3 ng/ μ l (9 ng/reaction). Initiate reaction by adding 3 μ l of diluted **BRDT** to wells designated for the “Negative Control” “Positive Control”, and “Test Inhibitor”. Discard any remaining diluted BRD protein after use.
- 9) Incubate at room temperature for 2 hours.
- 10) Read the fluorescent intensity in a microtiter-plate reader capable of TR-FRET.

Instrument Settings

| | |
|-----------------------|-----------------|
| Reading Mode | Time Resolved |
| Excitation Wavelength | 340 \pm 20 nm |
| Emission Wavelength | 620 \pm 10 nm |
| Lag Time | 60 μ s |
| Integration Time | 500 μ s |
| Excitation Wavelength | 340 \pm 20 nm |
| Emission Wavelength | 665 \pm 10 nm |
| Lag Time | 60 μ s |
| Integration Time | 500 μ s |

CALCULATING RESULTS:

Two sequential measurements should be conducted. Tb-donor emission should be measured at 620 nm followed by dye-acceptor emission at 665 nm. Data analysis is performed using the TR-FRET ratio (665 nm emission/620 nm emission).

When percentage activity is calculated, the FRET value from the negative control can be set as zero percent activity and the FRET value from the positive control can be set as one hundred percent activity.

$$\% \text{ Activity} = \frac{\text{FRET}_s - \text{FRET}_{neg}}{\text{FRET}_p - \text{FRET}_{neg}} \times 100\%$$

Where FRET_s = Sample FRET, FRET_{neg} = Negative control FRET, and FRET_p = Positive control FRET.


OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com

EXAMPLE OF ASSAY RESULTS:



Inhibition of BRDT (BD1 + BD2) by (+)-JQ1 (BPS Cat. #27401), measured using the *BRDT TR-FRET Assay Kit*, BPS Bioscience # 32616. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

RELATED PRODUCTS:

| <u>Product Name</u> | <u>Catalog</u> | <u>Size</u> |
|-------------------------------------|----------------|-------------|
| BET Bromodomain Ligand | 33000 | 0.5 mL |
| Bromodomain Non-acetylated Ligand 1 | 33005 | 0.5 mL |
| BRDT (22-138), His-tag* | 31101 | 100 µg |
| BRDT (257-382), His-tag* | 31100 | 100 µg |
| BRDT, BD1 and BD2 (22-382), GST-tag | 31061 | 100 µg |
| BRD1 (561 – 668), His-tag* | 31010 | 100 µg |
| BRD2 (65 – 459), His-tag* | 31025 | 100 µg |
| BRD3 (29 – 417), His-tag* | 31034 | 100 µg |
| BRD4 (49 – 460), His-tag* | 31045 | 100 µg |
| BRD1 Inhibitor Screening Kit | 32521 | 384 rxns. |
| BRD2 (BD2) Inhibitor Screening Kit | 32522 | 384 rxns. |
| BRD3 (BD1) Inhibitor Screening Kit | 32513 | 384 rxns. |
| BRD3 (BD2) Inhibitor Screening Kit | 32523 | 384 rxns. |
| BRD4 (BD1) Inhibitor Screening Kit | 32514 | 384 rxns. |
| BRD4 (BD2) Inhibitor Screening Kit | 32524 | 384 rxns. |
| (+)-JQ1 Inhibitor | 27401 | 1 mg |

*also available as GST-tag

Note: Tb-labeled donor and dye-labeled acceptor are products of Cisbio Bioassays.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com