



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Data Sheet

MTAP

human, recombinant, N-terminal GST tag
 Catalog # 50305
 Lot#: 80902 Conc.: 2.5 mg/ml

Formulated in: 45 mM Tris-HCl, pH 8.0,
 124 mM NaCl, 2.4 mM KCl, 18 mM
 glutathione, 10% glycerol, and 3 mM DTT.

Stability: >6 months at -80°C

References:


1. Fernandez-Irigoyen, J. *et al.*, *Biochem. J.* **411** (2), 457-465 (2008)
2. Hellerbrand, C. *et al.*, *Carcinogenesis* **27** (1), 64-72 (2006)
3. Ong, S-E., *et al.* *PNAS* **106**(12): 4617-4622 (2009)

Description: Human MTAP (GenBank Accession No. NM_002451), amino acids 2-end, with N-terminal GST tag, MW=57 kDa, expressed in an *E. coli* expression system.

Assay Conditions: MTAP activity was measured using a xanthine oxidase-coupled method. MTA cleavage by MTAP produce adenine, which is then converted to 2,8-dihydroxyadenine by xanthine oxidase, leading to an increase in absorbance at 305 nm. 50 uL reaction mixtures contained MTAP, 50 mM KPO₄, pH 7.4, 200 uM MTA, 0.02 units Xanthine Oxidase, and 1 mM DTT. Activity was measured at 37°C by continuous monitoring the absorbance at 305 nm for 30 min.

Application: Suitable for protein binding studies, Western blotting, or as an immunogen.

Quality Assurance



**10% SDS-PAGE
 Coomassie staining**

Lane 1:
 4 µg MTAP
Lane 2:
 Protein Marker

MW: 57 kDa
Purity: ≥80%

