



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet

JMJD2C Chemiluminescent Assay Kit

Catalog # 50405

DESCRIPTION: The *JMJD2C Chemiluminescent Assay Kit* is designed to measure JMJD2C activity for screening and profiling applications. JMJD2C is a JmjC-domain protein that exhibits demethylation activity toward H3-K9Me3 and H3-K36Me3. The *JMJD2C Chemiluminescent Assay Kit* comes in a convenient format, with a 96-well plate precoated with the methylated histone H3 peptide substrate, primary antibody, the secondary HRP-labeled antibody, demethylase assay buffer, and purified JMJD2C for 100 enzyme reactions. The key to the *JMJD2C Chemiluminescent Assay Kit* is a highly specific antibody that recognizes demethylated substrate. With this kit, only three simple steps on a microtiter plate are required for methyltransferase detection. First, a sample containing JMJD2C enzyme is incubated with a sample containing assay buffer for one hour. Next, primary antibody is added. Finally, the plate is treated with an HRP-labeled secondary antibody followed by the addition of the HRP substrate to produce chemiluminescence that can then be measured using a chemiluminescence reader.

COMPONENTS:

Catalog #	Component	Amount	Storage	
50105	JMJD2C (GASC1, KDM4C)	20 µg	-80°C	Avoid freeze/ thaw cycles!
52140E	Primary antibody 5	12.5 µl	-80°C	
52130H	Secondary HRP-labeled antibody 1	10 µl	-80°C	
	4x JMJD2C Direct Assay Buffer	3 x 1 ml	-80°C	
52100	Blocking buffer 4	50 ml	+4°C	
	HRP chemiluminescent substrate A	6 ml	+4°C	
	HRP chemiluminescent substrate B	6 ml	+4°C	
	White microplate precoated with histone substrate	1	+4°C	

MATERIALS REQUIRED BUT NOT SUPPLIED:

TBST buffer (1 x TBS, pH 8.0, containing 0.05% Tween20)
Luminometer or fluorescent microplate reader capable of reading chemiluminescence
Adjustable micropipettor and sterile tips
Rotating or rocker platform

APPLICATIONS: Great for studying enzyme kinetics and HTS applications.

*“*Note: The buffer in this kit was reformulated in May of 2015 with a different reducing agent to improve assay performance. The old formulation can still be purchased upon special request.”*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

CONTRAINDICATIONS: DMSO >1%, strong acids or bases, ionic detergents, high salt

STABILITY: One year from date of receipt when stored as directed.

REFERENCE: Whetstine JR ea. *Cell* 2006; **125**: 467.

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

Step 1:

- 1) Rehydrate the microwells by adding 200 μ l of TBST buffer (1 x TBS, pH 8.0, containing 0.05% Tween-20) to every well. Incubate 15 minutes at room temperature. Tap the strip plate onto clean paper towels to remove liquid.
- 2) Prepare master mix: N wells x (7.5 μ l **4x JMJD2C Direct Assay Buffer 1** + 12.5 μ l distilled water). Add 25 μ l of master mixture to each well.
- 3) Add 10 μ l of inhibitor solution of each well designated "Test Inhibitor". For the "Positive Control" and "Blank" add 10 μ l of the same solution without inhibitor (Inhibitor buffer). *Note: Keep final DMSO concentration \leq 1%.*

	Blank	Positive Control	Test Inhibitor
4x JMJD2C direct assay buffer 1	7.5 μ l	7.5 μ l	7.5 μ l
Distilled water	12.5 μ l	12.5 μ l	12.5 μ l
Test Inhibitor/Activator	-	-	10 μ l
Inhibitor buffer (no inhibitor)	10 μ l	10 μ l	-
1x JMJD2C direct assay buffer	20 μ l	-	-
JMJD2C (10 ng/ μ l)	-	20 μ l	20 μ l
Total	50 μl	50 μl	50 μl

- 4) Dilute 1 part **4x JMJD2C Direct Assay Buffer 1** with 3 parts distilled water (4-fold dilution) to make **1x JMJD2C Direct Assay Buffer**. Make only a sufficient quantity needed for the assay; store remaining stock solution in aliquots at -20°C.
- 5) Add 20 μ l of **1x JMJD2C Direct Assay Buffer 1** to wells designated as "Blank".
- 6) Thaw **JMJD2C** on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Aliquot **JMJD2C** enzyme into single use aliquots. Store remaining

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

undiluted enzyme in aliquots at -80°C. *Note: **JMJD2C** is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.*

- 7) Dilute **JMJD2C** in **1x JMJD2C Direct Assay Buffer** at 10 ng/μl. Keep diluted enzyme on ice until use. Discard any unused diluted enzyme after use.
- 8) Initiate reaction by adding 20 μl of diluted **JMJD2C** prepared as described above to wells designated "Positive Control" and "Test Inhibitor". Incubate at room temperature for one hour.
- 9) Wash the strip plate three times with TBST buffer. Blot dry onto clean paper towels.
- 10) Add 100 μl of **Blocking buffer 4** to every well. Shake on a rotating platform for 10 minutes. Remove the supernatant from the wells.

Step 2:

- 1) Dilute "**Primary antibody 5**" 800-fold with **Blocking buffer 4**.
- 2) Add 100 μl per well. Incubate 1 hour at room temperature with slow shaking.
- 3) Wash strip plate with TBST buffer and incubate in **Blocking buffer 4** as described in steps 1-9 and 1-10.

Step 3:

- 1) Dilute "**Secondary HRP-labeled antibody 1**" 1,000-fold with **Blocking Buffer 4**.
- 2) Add 100 μl per well. Incubate for 30 min. at room temperature with slow shaking.
- 3) Wash strip plate with TBST buffer and incubate in **Blocking buffer 4** as described in step 1-9 and 1-10.
- 4) Just before use, mix on ice 50 μl **HRP chemiluminescent substrate A** and 50 μl **HRP chemiluminescent substrate B** and add 100 μl per well. Discard any unused chemiluminescent reagent after use.
- 5) Immediately read sample in a luminometer or microtiter-plate capable of reading chemiluminescence.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com




6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Reading Chemiluminescence:

Chemiluminescence is the emission of light (luminescence) which results from a chemical reaction. The detection of chemiluminescence requires no wavelength selection because the method used is emission photometry and is not emission spectrophotometry.

To properly read chemiluminescence, make sure the plate reader is set for LUMINESCENCE mode. Typical integration time is 1 second, delay after plate movement is 100 msec. Do not use a filter when measuring light emission. Typical settings for the Synergy 2 BioTek plate reader are: use the "hole" position on the filter wheel; Optics position: Top; Read type: endpoint. Sensitivity may be adjusted based on the luminescence of a control assay without enzyme (typically we set this value as 100).

Example of Assay Results:



JMJD2C enzyme activity, measured using the *JMJD2C Chemiluminescent Assay Kit*, BPS Bioscience #50405. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

RELATED PRODUCTS

<u>Product Name</u>	<u>Catalog #</u>	<u>Size</u>
JMJD2A recombinant protein	50103	20 µg
JMJD2B recombinant protein	50104	20 µg
JMJD2C recombinant protein	50105	20 µg
JMJD3 recombinant protein	50115	20 µg
JMJD2A Homogeneous Assay Kit	50413	384 reactions
JMJD2B Homogeneous Assay Kit	50414	384 reactions
JMJD2C Homogeneous Assay Kit	50415	384 reactions
LSD1 Assay Kit (96 well)	50106	100 reactions
LSD1 recombinant protein	50100	50 µg
LSD1 substrate	50101	500 µl
COREST recombinant protein	50274	100 µg

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
 San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

TROUBLESHOOTING GUIDE

Problem	Possible Cause	Solution
Luminescence signal of positive control reaction is same as "blank" value.	JMJD2C has lost activity	Enzyme loses activity upon repeated freeze/thaw cycles. Use fresh JMJD2C, BPS Bioscience #50105. Store enzyme in single-use aliquots. Increase time of enzyme incubation. Increase enzyme concentration.
	Antibody reaction is insufficient	Increase time for primary antibody incubation. Avoid freeze/thaw cycles of antibodies.
	Incorrect settings on instruments	Refer to instrument instructions for settings to increase sensitivity of light detection.
	Chemiluminescent reagents mixed too soon	Chemiluminescent solution should be used within 15 minutes of mixing. Ensure both reagents are properly mixed.
Luminescent signal is erratic or varies widely among wells	Inaccurate pipetting/technique	Run duplicates of all reactions. Use a multichannel pipettor. Use master mixes to minimize errors.
	Bubbles in wells	Pipette slowly to avoid bubble formation. Tap plate lightly to disperse bubbles; be careful not to splash between wells.
Background (signal to noise ratio) is high	Insufficient washes	Increase number of washes. Increase wash volume. Increase Tween-20 concentration to 0.1% in TBST.
	Sample solvent is inhibiting the enzyme	Run negative control assay including solvent. Maintain DMSO level at <1%. Increase time of enzyme incubation.
	Results are outside the linear range of the assay	Use different concentrations of JMJD2C, BPS Bioscience #50105 to create a standard curve.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.
 To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**
 Or you can Email us at: info@bpsbioscience.com
 Please visit our website at: www.bpsbioscience.com