



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Data Sheet

EZH2 A738T /EED/SUZ12/RbAp48/AEBP2

Human, Recombinant, N-terminal FLAG-Tag (EED), N-Terminal His-Tag (Rest)

Catalog #: 51061

Lot #: 161118-1 **Conc.:** 0.83 mg/ml

Formulated in: 40 mM Tris-HCl, pH 8.0, 110 mM NaCl, 2.2 mM KCl, 20% glycerol

Stability: At least 6 months at -80°C . Avoid freeze/thaw cycles. Protein may be diluted to $\geq 100 \mu\text{g/ml}$ in PBS + glycerol and stored at -80°C .

References:

- Varambally, S. et al. Science 322 (5908), 1695-1699 (2008).
- Rakotobe, D. et al., Virol. J. 5, 32 (2008).


Description: Mutant version of our regular EZH2 5-member complex (Cat. # 51004), but with an Ala-to-Thr mutation on a.a. 738 of the EZH2 protein. Complex of human EZH2 (GenBank Accession No. NM_004456), (a-a 2-end ; A738T*) with N-terminal His tag, MW= 86 kDa, human EED (NM_003797) (a-a 2-end) with N-terminal Flag tag, MW= 51 kDa, human SUZ12 (NM_015355) (a-a 2-end) with N-terminal His tag, MW = 87 kDa, Human AEBP2 (NM_153207) (a-a 2-end) with N-terminal His tag, MW= 53 kDa, and human RbAp48 (NM_005610) (a-a 2-end) with N-terminal His tag, MW = 48 kDa, co-expressed in baculovirus expression system.

Specific Activity: 1.33 pmol/min/ μg
 50 μl reaction mix (20 mM phosphate buffer pH 7.4, 0.05% Tween-20, 10 μM S-adenosylmethionine, and various amount of enzyme complex) add to the wells coated with the substrate. Incubate for 1 hr. Add antibody against methylated K27 residue of histone H3, incubate 1 hr. Then, add secondary HRP-labeled antibody and incubate 30 min. Finally, add HRP chemiluminescent substrates and read luminescence.

Application: Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.

Quality Assurance

Specific Activity



6-20% SDS-PAGE Coomassie

Lane 1:

EZH2(A738T)-5m

Lane 2:

Protein Marker

Purity: $\geq 90\%$


MW: EZH2(Mut) 86 kDa

SUZ12 87 kDa

AEBP 53 kDa

EED 51 kDa

RbAp48 48 kDa



OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone 1.858.829.3082 Fax 1.858.481.8694

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com