



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet
PRMT1 Homogeneous Assay Kit
Catalog #52054

DESCRIPTION: The *PRMT1 Homogeneous Assay Kit* is designed to measure PRMT1 activity for screening and profiling applications. PRMT1 is a histone methyltransferase that exhibits methylation activity toward H4-R3. The *PRMT1 Homogeneous Assay Kit* comes in a convenient AlphaLISA® format, with a 384-well plate, biotinylated histone H4 peptide substrate, primary antibody, methylation assay buffer, and purified PRMT1 for 384 enzyme reactions. The key to the *PRMT1 Homogeneous Assay Kit* is a highly specific antibody that recognizes methylated histone substrate. With this kit, only three simple steps on a microtiter plate are required for methyltransferase detection. First, a sample containing PRMT1 enzyme is incubated with the biotinylated substrate for one hour. Next, acceptor beads and primary antibody are added, then donor beads, followed by reading the Alpha-counts.

COMPONENTS:

| Cat. # | Components | Amount | Storage | |
|---------|---|------------|---------|---|
| 51040 | PRMT1 | 1 µg | -80°C | (Avoid freeze/ thaw cycles!) |
| 52120 | 20 µM S-adenosylmethionine | 500 µl | -80°C | |
| 52150-3 | Primary antibody 4-3 | 4 x 100 µl | -80°C | |
| | Biotinylated histone H4 peptide substrate | 500 rxns | -80°C | |
| 52170-A | 4x HMT assay Buffer 2A | 3 ml | -20°C | |
| | 4x Detection Buffer | 2 ml | -20°C | |

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

AlphaLISA anti-rIgG acceptor beads, 5 mg/ml (Perkin Elmer #AL104C)
AlphaScreen Streptavidin-conjugated donor beads, 5 mg/ml (Perkin Elmer # 6760002)
OptiPlate-384 (Perkin Elmer #6007290)
AlphaScreen microplate reader

APPLICATIONS: Great for studying enzyme kinetics and HTS applications.

CONTRAINDICATIONS: Green and blue dyes that absorb light in the AlphaScreen signal emission range (520-620 nm), such as Trypan Blue. Avoid the use of the potent singlet oxygen quenchers such as sodium azide (NaN₃) or metal ions (Fe²⁺, Fe³⁺, Cu²⁺, Zn²⁺ and Ni²⁺). The presence of >1% RPMI 1640 culture medium leads to a signal reduction due to the presence of excess biotin and iron in this medium. MEM, which lacks these components, does not affect AlphaScreen assays.

STABILITY: One year from date of receipt when stored as directed.

REFERENCE(S): Li Y., et al., J. Neurosci. 2015; **35(37)**: 12890-12902.

FOR RESEARCH USE ONLY. NOT FOR HUMAN, DIAGNOSTIC, OR THERAPEUTIC USE.

To place your order, please contact us by Phone: **1.858.202.1401** Fax: **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
 San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

Step 1:

- 1) Re-suspend tube with **Biotinylated histone H4 peptide substrate** in 500 µl of distilled water.
- 2) Prepare the master mixture: N wells x (2.5 µl **4x HMT Buffer 2A** + 1 µl **S-adenosyl-methionine** (20 µM) + 1.0 µl **Biotinylated substrate**). Add 4.5 µl to wells designated "Positive Control", "Test Sample", and "Blank". To wells labeled "Substrate Control", add 2.5 µl **4x HMT Buffer 2A** + 1.0 µl **Biotinylated substrate** + 1.0 µl water.
- 3) Add 2.5 µl of Inhibitor solution of each well labeled as "Test Inhibitor". For the "Positive Control", "Substrate Control" and "Blank", add 2.5 µl of the same solution without inhibitor (Inhibitor buffer).
- 4) Prepare **1x HMT Buffer 2A** by adding 1 part of **4x HMT Buffer 2A** to 3 parts water (v/v).
- 5) Thaw **PRMT1** on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Aliquot **PRMT1** enzyme into single use aliquots. Store remaining undiluted enzyme in aliquots at -80°C. *Note: PRMT1 is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.*
- 6) Dilute **PRMT1** in **1X HMT assay Buffer 2A** at 0.5 ng/µl. Keep diluted enzyme on ice until use. Discard any unused diluted enzyme after use.

| | Blank | Positive Control | Test Sample | Substrate Control |
|---------------------------------|--------------|-------------------------|--------------------|--------------------------|
| 4x HMT assay Buffer 2A | 2.5 µl | 2.5 µl | 2.5 µl | 2.5 µl |
| 20 µM S-adenosylmethionine | 1 µl | 1 µl | 1 µl | - |
| Biotinylated substrate | 1 µl | 1 µl | 1 µl | 1 µl |
| H ₂ O | 3 µl | - | - | 1 µl |
| Test Inhibitor/Activator | - | - | 2.5 µl | - |
| Inhibitor Buffer (no inhibitor) | 2.5 µl | 2.5 µl | - | 2.5 µl |
| PRMT1 (0.5 ng/µl) | - | 3 µl | 3 µl | 3 µl |
| Total | 10 µl | 10 µl | 10 µl | 10 µl |

- 7) To the wells designated as "Blank", add 3 µl of water.
- 8) Initiate reaction by adding 3 µl of diluted **PRMT1** enzyme to the wells designated "Positive Control", "Substrate Control", and "Test Sample". Incubate at room temperature for 1 hour.

FOR RESEARCH USE ONLY. NOT FOR HUMAN, DIAGNOSTIC, OR THERAPEUTIC USE.

To place your order, please contact us by Phone: **1.858.202.1401** Fax: **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com


Step 2:

Note: Protect your samples from direct exposure to light!

- 1) Prepare **1x Detection Buffer** by adding 1 part of **4x Detection Buffer** to 3 parts water (v/v).
- 2) Dilute anti-Rabbit Acceptor beads (Perkin Elmer #AL104C) 1:250-fold with **1x Detection Buffer**. Add 5 μ l per well. Shake plate briefly.
- 3) Dilute **Primary antibody 4-3** 10-fold with **1x Detection Buffer**. Add 5 μ l per well. Shake plate. Incubate 30 min. at room temperature.

Step 3:

- 1) Dilute Streptavidin-conjugated donor beads (PE #6760002) 125-fold with **1x Detection Buffer**. Add 10 μ l per well. Incubate for 10 min. at room temperature.
- 2) Read Alpha-counts. The "Blank" value is subtracted from all other values.

Example of Assay Results:

PRMT1 enzyme activity, measured using the *PRMT1 Homogeneous Assay Kit*, BPS Bioscience Cat. #52054. Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com

FOR RESEARCH USE ONLY. NOT FOR HUMAN, DIAGNOSTIC, OR THERAPEUTIC USE.

To place your order, please contact us by Phone: **1.858.202.1401** Fax: **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

RELATED PRODUCTS:

| | | |
|--|---------|---------------|
| PRMT8 Homogeneous Assay Kit | #52058 | 384 reactions |
| PRMT3 Homogeneous Assay Kit | #52055 | 384 reactions |
| PRMT5 Homogeneous Assay Kit | #52052 | 384 reactions |
| PRMT6 Homogeneous Assay Kit | #52056 | 384 reactions |
| PRMT1 Chemiluminescent Assay Kit | #52004L | 96 reactions |
| PRMT3 Chemiluminescent Assay Kit | #52005L | 96 reactions |
| PRMT4 Chemiluminescent Assay Kit | #52041L | 96 reactions |
| PRMT5 Chemiluminescent Assay Kit | #52002L | 96 reactions |
| PRMT6 Chemiluminescent Assay Kit | #52046 | 96 reactions |
| PRMT1 recombinant protein (<i>E. coli</i>) | #51040 | 50 µg |
| PRMT1 recombinant protein (Sf9) | #51041 | 20 µg |
| PRMT3 recombinant protein | #51043 | 50 µg |
| PRMT4 (CARM 1) recombinant protein | #51047 | 20 µg |
| PRMT5 recombinant protein (HEK293) | #51045 | 20 µg |
| PRMT5/MEP50 recombinant protein (Sf9) | #51048 | 20 µg |
| PRMT6 recombinant protein | #51049 | 20 µg |
| PRMT7 recombinant protein | #51054 | 20 µg |
| PRMT8 recombinant protein | #51052 | 20 µg |
| PRMT9 recombinant protein | #51053 | 20 µg |

AlphaScreen® and AlphaLISA® are registered trademarks of PerkinElmer, Inc.

FOR RESEARCH USE ONLY. NOT FOR HUMAN, DIAGNOSTIC, OR THERAPEUTIC USE.

To place your order, please contact us by Phone: **1.858.202.1401** Fax: **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com