



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Ct. West, Ste. B
San Diego, CA 92121
Tel: 1.858.829.3082
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet
TDO Fluorogenic Inhibitor Screening Assay Kit
(384 Well Format)
Catalog # 72049

DESCRIPTION: The *TDO Fluorogenic Inhibitor Screening Assay Kit* is designed to measure enzyme inhibition of tryptophan 2,3-dioxygenase (TDO). The kit comes in a convenient format, with enough reaction solution and enzyme to perform a total of 400 reactions. The *TDO Fluorogenic Inhibitor Screening Assay Kit* is simple to use and detects fluorescence at long wavelengths, which minimizes potential errors due to compound interference. In the assay, the inhibitor and enzyme are added to a sample containing L-Trp substrate. After a 1 hour incubation at room temperature, the fluorescence solution is added and incubated at 37°C for four hours. Activity is measured by reading sample fluorescence at $\lambda=510$ nm following excitation of the reaction product at $\lambda=400$ nm.

BACKGROUND: L-tryptophan (L-Trp) is an essential amino acid necessary for protein synthesis in mammalian cells and the L-Trp to kynurenine (Kyn) pathway is firmly established as a key regulator of innate and adaptive immunity. Catabolism of L-Trp to Kyn maintains an immunosuppressive microenvironment by starving immune cells of L-Trp and releasing degradation products of L-Trp that have immunosuppressive functions. Tryptophan 2,3 dioxygenase (TDO), a rate limiting enzymes in this pathway, is upregulated in many tumors, providing cancer cells with an avenue for immune evasion.

COMPONENTS:

Catalog #	Component	Amount	Storage	
71195	TDO His-Tag	90 μ g	-80°C	(Avoid freeze/ thaw cycles!)
73010	TDO Fluorogenic Reaction Solution	3 x 10 ml	-80°C	
73006	TDO Buffer	3 ml	-80°C	
	Fluorescence Solution	4 x 1 ml	-80°C	
	Black 384 Well Assay-Plate	1		
	Plate sealing film	1		

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Fluorimeter capable of excitation at 390-410 nm and detection at 500-520 nm
Adjustable micropipettor and sterile tips
Rotating or rocker platform

APPLICATIONS: Useful for the study of TDO enzymology, inhibitor screening, and selectivity profiling.

STABILITY: At least one year from date of receipt when stored as directed.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082**, Fax **1.858.481.8694**
Or you can Email us at: info@bpsbioscience.com
Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Ct. West, Ste. B
San Diego, CA 92121
Tel: 1.858.829.3082
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

CONTRAINDICATIONS: DMSO >0.5%, strong acids or bases, ionic detergents, high salt.
Warning: the Fluorescence Solution contains a component that is known to be a skin and eye irritant. Use caution and appropriate personal protective equipment when handling this component.

REFERENCE: Seegers, N., *et al. J. Biomol. Screen.* 2014. **19(9):**1266-74.

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate. Use slow shaking for all incubations.

Step 1:

- 1) Thaw reaction solution and aliquot 72 μ l into each well. Note: TDO Reaction Solution may contain a precipitate after thawing. Please ensure the mixture is fully solubilized by gentle mixing before aliquoting. Do not vortex.
- 2) Add 4 μ l of inhibitor solution (no more than 10% DMSO) to each well designated "Test Inhibitor". For the wells designated "Positive Control" and "Blank", add 4 μ l of the same solution without inhibitor (inhibitor buffer). Note: Keep the final DMSO concentration below 0.5%.
- 3) Dilute **TDO** in **TDO Buffer** at 50 ng/ μ l. Keep diluted protein on ice until use. Discard any unused diluted protein after use.

	Blank	Positive Control	Test Inhibitor
Reaction Solution	72 μ l	72 μ l	72 μ l
Test Inhibitor	-	-	4 μ l
Inhibitor buffer (no inhibitor)	4 μ l	4 μ l	-
TDO Buffer	4 μ l	-	-
TDO (50 ng/ μ l)	-	4 μ l	4 μ l
Total	80 μl	80 μl	80 μl

- 4) Add 4 μ l of **TDO Assay Buffer** to the wells designated "Blank".
- 5) Initiate reaction by adding 4 μ l of diluted **TDO** prepared as described above to the wells labeled "Positive Control", and "Test Inhibitor". Incubate at room temperature for 1 hour.
- 6) Add 8 μ l **Fluorescence Solution** to each well. Seal the plate and incubate at 37°C for four hours. Allow plate to cool for 10 minutes.


OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082**, Fax **1.858.481.8694**
Or you can Email us at: info@bpsbioscience.com
Please visit our website at: www.bpsbioscience.com

- 7) Unseal the plate and measure fluorescence in a fluorimeter capable of excitation at 400 nm and emission at 510 nm. Subtract "Blank" value from all other values.

EXAMPLE OF ASSAY RESULTS:

**TDO2 Inhibition by 680C91
 Fluorescence Assay (384 well format)**



Inhibition of TDO activity by 680C91, measured using the *TDO Fluorogenic Inhibitor Screening Assay Kit (384 well format)*, BPS Bioscience, Catalog # 72049. Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com.

RELATED PRODUCTS:

Product Name	Catalog #	Size
IDO1, His-tag	71182	50 μg
IDO2, His-tag	71194	50 μg
TDO, His-tag	71195	50 μg
IDO1 Fluorogenic Inhibitor Screening Kit	72034	96 rxns
IDO1 Inhibitor Screening Assay Kit	72021	96 rxns
IDO2 Inhibitor Screening Assay Kit	72022	96 rxns
TDO Inhibitor Screening Assay Kit	72023	96 rxns
IDO1 Cell-Based Assay Kit	72031	100 rxns
TDO Cell-Based Assay Kit	72033	100 rxns
TDO-HEK293 Recombinant Cell line	60534	2 vials
TDO Cellular Activity QuickDetect™ Supplements	62002-1	100 rxns
N-formylkynurenine	73000	2 mg
NLG919	27337-1	10 mg
INCB024360	27338-1	10 mg
TDO Reaction Solution	73005	1 ml
TDO Assay Buffer	73006	1 ml

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082**, Fax **1.858.481.8694**
 Or you can Email us at: info@bpsbioscience.com
 Please visit our website at: www.bpsbioscience.com