



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet
HER4 Assay Kit
Catalog # 78005
96 Reactions

BACKGROUND: Human HER4 kinase, also known as Erb-B4, Proto-Oncogene-Like Protein C-ErbB-4, and Tyrosine Kinase-Type Cell Surface Receptor when mutated has been implicated in cancer. There has also been connection to schizophrenia with single-nucleotide polymorphisms.

DESCRIPTION: The *HER4 Assay Kit* is designed to measure HER4 activity for screening and profiling applications using Kinase-Glo[®] MAX as a detection reagent. The *HER4 Assay Kit* comes in a convenient 96-well format, with enough purified HER4, Protein Tyrosine Kinase Substrate (Poly-Glu,Tyr 4:1), ATP, and kinase assay buffer for 96 enzyme reactions.

COMPONENTS:

Catalog #	Reagent	Amount	Storage	
40232	HER4, GST-Tag	30 µg	-80°C	Avoid multiple freeze/thaw cycles!
79334	5x Kinase assay buffer	1.5 ml	-20°C	
79686	ATP (500 µM)	100 µl	-20°C	
40217	Protein Tyrosine Kinase Substrate (Poly-Glu,Tyr 4:1) (10 mg/ml)	100 µl	-20°C	
79696	96-well plate, white	1	RT	

APPLICATIONS: Useful for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: Up to 6 months when stored as recommended.

REFERENCE(S):

1. Plowman, Gregory D., et al. "Ligand-specific activation of HER4/p180erbB4, a fourth member of the epidermal growth factor receptor family." *Proceedings of the National Academy of Sciences* 90.5 (1993): 1746-1750.
2. Elenius, Klaus, et al. "Activation of HER4 by heparin-binding EGF-like growth factor stimulates chemotaxis but not proliferation." *The EMBO journal* 16.6 (1997): 1268-1278.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Kinase-Glo MAX (Promega #V6071)
Dithiothreitol (DTT, 0.5 M; optional)
Microplate reader capable of reading luminescence
Adjustable micropipettor and sterile tips
30°C incubator

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

Thaw **5x Kinase assay buffer**, **ATP (500 µM)**, and **Protein Tyrosine Kinase Substrate (Poly-Glu,Tyr 4:1)**. (Optional: If desired, add 30 µl of 0.5 M DTT to **5x Kinase assay buffer**).

- 1) Prepare the master mixture (25 µl per well): N wells x (10 µl **5x Kinase assay buffer** + 1 µl **ATP (500 µM)** + 1 µl **Protein Tyrosine Kinase Substrate (Poly-Glu,Tyr 4:1)** + 13 µl distilled water). Add 25 µl to every well.

	Positive Control	Test Inhibitor	Blank
5x Kinase assay buffer	10 µl	10 µl	10 µl
ATP (500 µM)	1 µl	1 µl	1 µl
Poly-Glu,Tyr (10 mg/ml)	1 µl	1 µl	1 µl
Water	13 µl	13 µl	13 µl
Test Inhibitor	-	5 µl	-
10% DMSO in water (Inhibitor buffer)	5 µl	-	5 µl
1x Kinase buffer	-	-	20 µl
HER4, GST-Tag (15 ng/µl)	20 µl	20 µl	-
Total	50 µl	50 µl	50 µl

- 2) Add 5 µl of Inhibitor solution of each well labeled as "Test Inhibitor." For the "Positive Control" and "Blank," add 5 µl of 10% DMSO in water (Inhibitor buffer). *Note: Final DMSO concentration must be ≤1%. Higher DMSO levels can significantly decrease the enzyme activity. For example, to test an inhibitor dissolved in 100% DMSO at 10 µM, dilute 1 mM inhibitor with water to make a 100 µM inhibitor in 10% DMSO(aq). Then, add 5 µl of the 100 µM solution into the 50 µl assay to make a 1% DMSO concentration in the final reaction mixture.*
- 3) Prepare 3 ml of **1x Kinase assay buffer** by mixing 600 µl of **5x Kinase assay buffer** with 2400 µl water. 3 ml of **1x Kinase assay buffer** is sufficient for 100 reactions.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

- 4) To the wells designated as "Blank," add 20 μ l of **1x Kinase assay buffer**.

- 5) Thaw **HER4, GST-Tag** on ice. Upon first thaw, briefly spin tube containing material to recover full content of the tube. Calculate the amount of **HER4, GST-Tag** required for the assay and dilute enzyme to 15 ng/ μ l with **1x Kinase assay buffer**. Store remaining undiluted material in aliquots at -80°C. *Note: HER4, GST-Tag is sensitive to freeze/thaw cycles. Avoid multiple freeze/thaw cycles. Do not re-use thawed aliquots or diluted material.*

- 6) Initiate reaction by adding 20 μ l of diluted **HER4, GST-Tag** to the wells designated "Positive Control" and "Test Inhibitor Control." Incubate at 30°C for 45 minutes.

- 7) Thaw Kinase-Glo Max reagent.

- 8) After the 45-minute reaction, add 50 μ l of Kinase-Glo Max reagent to each well. Cover plate with aluminum foil and incubate the plate at room temperature for 15 minutes.

- 9) Measure luminescence using the microplate reader. Value of "Blank" reading should be subtracted from all other measurements.


OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com

Example of Assay Results:



Inhibition of HER4, GST-Tag by Staurosporine, measured using the HER4 assay kit (BPS Bioscience #78005). *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

RELATED PRODUCTS:

<u>Product Name</u>	<u>Catalog #</u>	<u>Size</u>
EPHA2, His-tag	40190	10 μg
5x Kinase assay buffer	79334	10 ml
ATP (500 μM)	79686	200 μl
Protein Tyrosine Kinase Substrate (poly-Glu,Tyr 4:1)	40217	1 mg
EPHA1, GST-tag	40191	10 μg
EPHA3, GST-tag	40192	10 μg
EPHA4, GST-tag	40193	10 μg
EPHA6, GST-tag	40194	10 μg
EPHB2, His-tag	40200	10 μg
EPHB3, GST-tag	40186	10 μg
EPHB4, His-tag	40201	10 μg
EPHA6, GST-tag	40194	10 μg
EPHB1, GST-tag (Mouse)	40199	10 μg

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com