



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet
AKT1 Assay Kit
Catalog # 78038
96 Reactions

BACKGROUND: AKT1, also known as Protein Kinase B Alpha (PKB α), is a serine/threonine kinase found in various cell types throughout the body, where it plays a critical role in regulating cell survival, insulin signaling, angiogenesis and tumor formation as well as controlling apoptosis. Importantly, the signaling pathway involving AKT1 kinase appears to be essential for the normal development and function of the nervous system. Dysregulation of Akt1 contributes to a number of serious diseases, including cancer, diabetes, cardiovascular and neurological diseases.

DESCRIPTION: The *AKT1 Assay Kit* is designed to measure AKT1 activity for screening and profiling applications using Kinase-Glo[®] Kinase Assay as a detection reagent. The *AKT1 Assay Kit* comes in a convenient 96-well format, with enough purified recombinant AKT1 enzyme, AKT1 Substrate, ATP, and kinase assay buffer for 100 enzyme reactions.

COMPONENTS:

| Catalog # | Reagent | Amount | Storage | |
|-----------|--------------------------|-------------|------------|---|
| 40003 | AKT1 | 10 μ g | -80°C | Avoid multiple freeze/thaw cycles! |
| 79334 | 5x Kinase assay buffer 1 | 1.5 ml | -20°C | |
| 79686 | ATP (500 μ M) | 100 μ l | -20°C | |
| | AKT1 Substrate (5 mg/ml) | 100 μ l | -20°C | |
| 79696 | 96-well plate, white | 1 | Room Temp. | |

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Kinase-Glo[®] Kinase Assay (Promega #V6071)
Dithiothreitol (DTT, 0.5 M)
Microplate reader capable of reading luminescence
Adjustable micropipettor and sterile tips
30°C incubator

APPLICATIONS: Useful for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: Up to 6 months when stored as recommended.

REFERENCES:

1. Kuijl C, *et al.*, *Nature*. 2007 Nov 29; **450 (7170)**:725-30.
2. Vogiatzi, P. and Giordano, A. *Cancer Biol Ther*. 2007 Aug 3; **6(10)**:1521-1524.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

- 1) Thaw **5x Kinase assay buffer**, **ATP (500 μ M)**, and **AKT1 Substrate**.
- 2) Prepare the master mixture (25 μ l per well): N wells x (6 μ l **5x Kinase assay buffer** + 1 μ l **ATP (500 μ M)** + 1 μ l **AKT1 Substrate** + 17 μ l distilled water. Add 25 μ l to every well.

| | Positive Control | Test Inhibitor | Blank |
|---------------------------------|------------------|----------------|------------|
| 5x Kinase assay buffer | 6 μ l | 6 μ l | 6 μ l |
| ATP (500 μ M) | 1 μ l | 1 μ l | 1 μ l |
| AKT1 Substrate | 1 μ l | 1 μ l | 1 μ l |
| Water | 17 μ l | 17 μ l | 17 μ l |
| Test Inhibitor | - | 5 μ l | - |
| Inhibitor Buffer (no inhibitor) | 5 μ l | - | 5 μ l |
| 1x Kinase buffer | - | - | 20 μ l |
| AKT1 (5 ng/ μ l) | 20 μ l | 20 μ l | - |
| Total | 50 μ l | 50 μ l | 50 μ l |

- 1) Prepare 10X concentrated inhibitor in an aqueous-based solution. *Note: Final DMSO concentration must be \leq 1%. Higher DMSO levels can significantly decrease the enzyme activity. For example, to test an inhibitor at 10 μ M, dilute 1 mM inhibitor in 100% DMSO with water to make a 100 μ M inhibitor in 10% DMSO (aq). Then, add 5 μ l of the 100 μ M solution to the assay to make a 10 μ M in 1% DMSO concentration in the final reaction mixture.*
 - 3) Add 5 μ l of Inhibitor solution of each well labeled as "Test Inhibitor." For the "Positive Control" and "Blank," add 5 μ l of the same solution without inhibitor (Inhibitor buffer, i.e. 10% DMSO(aq)). Be sure to maintain the same concentration of DMSO in the Positive Control sample as the Test sample.
 - 2) Prepare 3 ml of **1x Kinase assay buffer** by mixing 600 μ l of **5x Kinase assay buffer** with 2,376 μ l distilled water and 24 μ l of 0.5M DTT. 3 ml of **1x Kinase assay buffer** is sufficient for 100 reactions.
 - 3) To the wells designated as "Blank," add 20 μ l of **1x Kinase assay buffer**.
 - 4) Thaw **AKT1** on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Calculate the amount of **AKT1** required for the assay and
- OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com


Please visit our website at: www.bpsbioscience.com

dilute enzyme to 5 ng/μl with **1x Kinase assay buffer**. Store remaining undiluted enzyme in aliquots at -80°C.

Note: AKT1 enzyme is sensitive to freeze/thaw cycles. Avoid multiple freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.

- 5) Initiate reaction by adding 20 μl of diluted **AKT1** to the wells designated "Positive Control" and "Test Inhibitor." Incubate at 30°C for 45 minutes.
- 6) Thaw Kinase-Glo Max reagent.
- 7) After the 45 minute reaction, add 50 μl of Kinase-Glo Max reagent to each well. Cover plate with aluminum foil and incubate the plate at room temperature for 15 minutes.
- 8) Measure luminescence using the microplate reader. Value of "Blank" reading should be subtracted from all other measurements.

Example of Assay Results:



Inhibition of AKT1 by A-674563, measured using the *AKT1 assay kit* (BPS Bioscience #78038). *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

RELATED PRODUCTS:

| <u>Product Name</u> | <u>Catalog #</u> | <u>Size</u> |
|----------------------------|-------------------------|--------------------|
| AKT1, His-tag | 40003 | 10 µg |
| 5x Kinase buffer 1 | 79334 | 10 ml |
| ATP (500 µM) | 79686 | 200 µl |
| AKT1, Inactive, His-tag | 40000 | 100 µg |
| AKT2, His-tag | 40011 | 10 µg |
| AKT2, Inactive, His-tag | 40001 | 100 µg |
| AKT3, His-tag | 40012 | 10 µg |

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com