



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court West, Suite B  
San Diego, CA 92121  
Tel: 1.858.202.1401  
Fax: 1.858.481.8694  
Email: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

## Data Sheet **RANK:RANKL TR-FRET Assay** Catalog # 79101

### DESCRIPTION:

The RANK:RANKL TR-FRET Assay is designed to measure the inhibition of RANK binding to RANKL in a homogeneous 384 reaction format. This FRET-based assay requires no time-consuming washing steps, making it especially suitable for high throughput screening applications. The assay procedure is straightforward and simple; a sample containing biotinylated RANK, RANKL, anti-His Tb donor, dye-labeled acceptor, and an inhibitor is incubated for two hours. Then, the fluorescence intensity is measured using a fluorescence reader.

### COMPONENTS:

Catalog #	Component	Amount	Storage	
70822	RANK, biotinylated	5 µg	-80°C	<b>(Avoid freeze/thaw cycles!)</b>
71051	RANKL-His	2 µg	-80°C	
	Anti-His Tb Donor	2 x 10 µl	-20°C	
	Dye-labeled Acceptor	2 x 10 µl	-20°C	
79311	3x Immuno Buffer 1	4 ml	-20°C	
	White, non-binding, low volume, 384-well microtiter plate	1	Room temp.	

### MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Fluorescence microplate reader capable of measuring Time Resolved Fluorescence Resonance Energy Transfer (TR-FRET)  
Adjustable micropipettor and sterile tips

**APPLICATIONS:** Great for screening small molecular inhibitors for drug discovery and HTS applications.

**STABILITY:** At least 6 months from date of receipt when stored as directed.

### REFERENCES:

1. Annis, A., *et al. J. Amer.n Chem. Soc.*, 2004 May; **126(4)**: 15495-15503.
2. Yan, T., *et al. J. Cell. Biochem.* 2001 Aug; **83(2)**: 320-325.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**  
Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)  
Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court West, Suite B  
San Diego, CA 92121  
Tel: 1.858.202.1401  
Fax: 1.858.481.8694  
Email: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

## ASSAY PROTOCOL:

All samples and controls should be tested in at least duplicate.

### Protocol for RANK assay

- 1) Dilute one part **3x Immuno Buffer 1** with 2 parts distilled water (3-fold dilution) to make **1x Immuno Buffer 1**. Make only a sufficient quantity needed for the assay; store remaining stock solution in aliquots at -20°C.
- 2) Dilute **Anti-His Tb Donor** 100-fold in **1x Immuno Buffer 1**. Make only sufficient quantities needed for the assay; store remaining stock solution in aliquots at -20°C.
- 3) Dilute **Dye-labeled Acceptor** 100-fold in **1x Immuno Buffer 1**. Make only sufficient quantities needed for the assay; store remaining stock solution in aliquots at -20°C.
- 4) Thaw **RANK, Biotinylated** on ice. Upon first thaw, briefly spin tube containing **RANK, Biotinylated** to recover the full contents of the tube. Aliquot into single-use aliquots. Store remaining undiluted **RANK** at -80°C immediately. *Note: **RANK, Biotinylated** is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted protein.*
- 5) Dilute **RANK, Biotinylated** in **1x Immuno Buffer 1** to 1.65 µg/ml. Make only sufficient quantities needed for the assay; store remaining stock solution in aliquots at -20°C.
- 6) Prepare the master mixture: N wells x (3 µl diluted **RANK, Biotinylated** + 5 µl diluted **Anti-His Tb Donor** + 5 µl diluted **Dye-labeled acceptor**). Add 13 µl to every well.
- 6) Add 2 µl of inhibitor solution to each well designated "Test Inhibitor." Add 2 µl of the same solution without inhibitor (inhibitor buffer) to the wells labeled "Negative Control" and "Positive Control."
- 7) Add 5 µl **1x Immuno Buffer 1** to wells designated for "Negative Control."

	Positive Control	Negative Control	Test Inhibitor
RANK, Biotinylated (diluted)	3 µl	3 µl	3 µl
Anti-His Tb Donor	5 µl	5 µl	5 µl
Dye-labeled Acceptor	5 µl	5 µl	5 µl
Test Inhibitor	-	-	2 µl
Inhibitor Buffer (no inhibitor)	2 µl	2 µl	-
1x Immuno Buffer 1	-	5 µl	-
RANKL-His (diluted)	5 µl	-	5 µl
<b>Total</b>	<b>20 µl</b>	<b>20 µl</b>	<b>20 µl</b>

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**  
Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)  
Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court West, Suite B  
San Diego, CA 92121  
Tel: 1.858.202.1401  
Fax: 1.858.481.8694  
Email: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

- 8) Thaw **RANKL-His** protein on ice. Upon first thaw, briefly spin tube containing protein to recover the full contents of the tube. Aliquot **RANKL-His** into single-use aliquots. Store remaining undiluted **RANKL-His** in aliquots at  $-80^{\circ}\text{C}$  immediately. *Note: **RANKL-His** is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted protein.*
- 9) Dilute **RANKL-His** in **1x Immuno Buffer 1** to  $0.5\ \mu\text{g/ml}$ . Initiate reaction by adding  $5\ \mu\text{l}$  of diluted **RANKL-His** to wells designated for the "Positive Control" and "Test Inhibitor." Discard any remaining diluted **RANKL-His** protein after use.
- 10) Incubate at room temperature for 1.5 hours.
- 11) Read the fluorescent intensity in a microtiter-plate reader capable of TR-FRET.

#### Instrument Settings

Reading Mode	Time Resolved
Excitation Wavelength	$320\pm 10\ \text{nm}$
Emission Wavelength	$620\pm 10\ \text{nm}$
Lag Time	$60\ \mu\text{s}$
Integration Time	$500\ \mu\text{s}$
Excitation Wavelength	$320\pm 20\ \text{nm}$
Emission Wavelength	$665\pm 10\ \text{nm}$
Lag Time	$60\ \mu\text{s}$
Integration Time	$500\ \mu\text{s}$

#### CALCULATING RESULTS:

Two sequential measurements should be conducted. Tb-donor emission should be measured at  $620\ \text{nm}$  followed by dye-acceptor emission at  $665\ \text{nm}$ . Data analysis is performed using the TR-FRET ratio ( $665\ \text{nm}$  emission/ $620\ \text{nm}$  emission).

If desired, data can be normalized to percent inhibition. Typically for inhibitor screens, the FRET value from the positive control is set to zero percent inhibition and the FRET value from the negative control is set to one hundred percent inhibition.


OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**  
Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)  
Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court West, Suite B  
San Diego, CA 92121  
Tel: 1.858.202.1401  
Fax: 1.858.481.8694  
Email: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

**EXAMPLE OF ASSAY RESULTS:**



**Figure Legend:** Inhibition of RANK:RANKL interaction with unlabeled RANK (BPS Bioscience, #70823). Data in the above graphs are expressed as FRET ratios. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at [info@bpsbioscience.com](mailto:info@bpsbioscience.com)*

**RELATED PRODUCTS:**

<u>Product</u>	<u>Catalog #</u>	<u>Size</u>
RANKL, His-Tag (Human)	71051	100 µg
RANK, Fc fusion (IgG1), Avi-tag (Human)	70823	100 µg
RANK, Fc fusion (IgG1), Biotin Labeled (Human)	70822	50 µg

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**  
Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)  
Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)