



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** info@bpsbioscience.com

## Data Sheet

### ***Mouse CTLA4[Biotin]:B7-1 Inhibitor Screening Assay Kit***

**Catalog # 79515**  
**Size: 96 reactions**

**DESCRIPTION:** The Mouse *CTLA4[biotin]:B7-1 Inhibitor Screening Assay Kit* is designed for screening and profiling inhibitors of mouse CTLA4:B7-1 signaling. This kit comes in a convenient 96-well format, with biotin-labeled mouse CTLA4, purified mouse B7-1, streptavidin-labeled HRP, and assay buffer for 100 binding reactions. The key to this kit is the high sensitivity of detection of biotin-labeled mouse CTLA4 by streptavidin-HRP. Only a few simple steps on a microtiter plate are required for the assay. First, B7-1 is coated on a 96-well plate. Next, CTLA4[biotin] is incubated with B7-1 on the plate. Finally, the plate is treated with streptavidin-HRP followed by addition of an HRP substrate to produce chemiluminescence, which can be measured using a chemiluminescence reader.

**BACKGROUND:** The activation of naïve T cells requires two signals, the specific T cell receptor recognition of MHC/Antigen on the surface of the antigen-presenting cell (APC), and the binding of B7-1 (CD80) ligand on the APC with the CD28 receptor on the T cell surface. Conversely, binding of CTLA4 to B7-1 on the T-cell surface results in an inhibitory signal and prevents T-cell activation. CTLA4:B7-1 interaction is an important drug target for the regulation of the host's response to cancer.

#### **COMPONENTS:**

Catalog #	Component	Amount	Storage	
79001	Mouse CTLA4 (CD152), Fc-Biotin-labeled	2 µg	-80°C	<b>(Avoid freeze/thaw cycles!)</b>
79058	Mouse B7-1 (CD86), Fc fusion	5 µg	-80°C	
	Streptavidin-HRP	15 µl	+4°C	
79311	3x Immuno Buffer 1	50 ml	-20°C	
79728	Blocking Buffer 2	50 ml	+4°C	
79670	ELISA ECL Substrate A	6 ml	+4°C	
79670	ELISA ECL Substrate B	6 ml	+4°C	
	White 96-well microplate	1	+4°C	

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

**MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:**

PBS (Phosphate buffered saline)

Luminometer or fluorescent microplate reader capable of reading chemiluminescence

Rotating or rocker platform

**APPLICATIONS:** This kit is useful for screening for inhibitors of mouse CTLA4 binding to B7-1.

**STABILITY:** One year from date of receipt when stored as directed.

**REFERENCES:**

1. Ohtani, H., *et al.*, *Lab Invest.* 1997; **77(3)**: 231-241.
2. Rovert, C., *et al.*, *N. Engl. J. Med.* 2011; **364**: 2517-25262.

**ASSAY PROTOCOL:**

All samples and controls should be tested in duplicate.

**Coating the plate with B7-1:**

- 1) Thaw **Mouse B7-1** on ice. Upon first thaw, briefly spin tube containing **B7-1** to recover the full contents of the tube. Aliquot into single use aliquots. Immediately store remaining B7-1 in aliquots at -80°C. *Note: B7-1 is very sensitive to freeze/thaw cycles. Avoid multiple freeze/thaw cycles.*
- 2) Dilute **B7-1** to 1 µg/ml in PBS.
- 3) Add 50 µl of diluted **B7-1** solution to each well and incubate overnight at 4°C. Leave a couple of wells empty (uncoated), for use with the "Ligand Control" (see below).
- 4) Dilute **3x Immuno Buffer 1** to **1x Immuno Buffer 1** in water. Dilute only enough required for the assay; store remaining 3x buffer at -20°C.
- 5) Decant to remove supernatant. Wash the plate 3 times with 100 µl **1x Immuno Buffer 1**. Tap plate onto clean paper towels to remove liquid.
- 6) Block wells by adding 100 µl of **Blocking Buffer 2** to each well. Incubate for 1 hour at room temperature. Remove supernatant as described in step 4.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
 San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** info@bpsbioscience.com

**Step 1:**

- 1) Prepare the master mixture: N wells × (10 µl **3x Immuno Buffer 1** + 15 µl H<sub>2</sub>O).
- 2) Add 25 µl of master mixture to each well. Use uncoated wells for the “Ligand Control”.
- 3) Add 5 µl of inhibitor solution to each well designated “Test Inhibitor”. For the “Positive Control”, “Ligand Control” and “Blank”, add 5 µl of the same solution without inhibitor (inhibitor buffer). Incubate at room temperature for one hour.

	<b>Blank</b>	<b>Ligand Control</b>	<b>Positive Control</b>	<b>Test Inhibitor</b>
3x Immuno Buffer 1	10 µl	10 µl	10 µl	10 µl
H <sub>2</sub> O	15 µl	15 µl	15 µl	15 µl
Test Inhibitor/Activator	–	–	–	5 µl
Inhibitor buffer (no inhibitor)	5 µl	5 µl	5 µl	–
1x Immuno Buffer 1	20 µl	–	–	–
CTLA4-biotin (0.05 µg/ml)	–	20 µl	20 µl	20 µl
<b>Total</b>	<b>50 µl</b>	<b>50 µl</b>	<b>50 µl</b>	<b>50 µl</b>

- 4) Thaw **Mouse CTLA4-biotin** on ice. Upon first thaw, briefly spin tube containing protein to recover full contents of the tube. Aliquot **CTLA4-biotin** into single use aliquots. Immediately store remaining undiluted protein in aliquots at -80°C. *Note: CTLA4-biotin is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted protein.*
- 5) Dilute **CTLA4-biotin** in **1x Immuno Buffer 1** to 0.05 µg/ml. Keep diluted protein on ice until use. Discard any unused diluted protein after use.
- 6) Add 20 µl of **1x Immuno Buffer 1** to the well designated “Blank”.
- 7) Initiate reaction by adding 20 µl of diluted **CTLA4-biotin** (see Step 1-5) to wells labeled “Positive Control”, “Ligand Control” and “Test Inhibitor”. Incubate at room temperature for two hours.
- 8) Decant to remove supernatant. Wash the plate 3 times with 100 µl/well **1x Immuno Buffer 1**. Tap plate onto clean paper towels to remove liquid.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**  
 Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)  
 Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

- 9) Block wells by adding 100  $\mu$ l of **Blocking Buffer 2** to each well. Incubate for 10 minutes at room temperature. Remove supernatant as in Step 1-8.

### **Step 2:**

- 1) Dilute **Streptavidin-HRP** 1000-fold with **Blocking Buffer 2**.
- 2) Add 100  $\mu$ l to each well. Incubate for 1 hour at room temperature with slow shaking.
- 3) Wash plate three times with **1x Immuno Buffer 1**. Tap onto clean paper towels to remove liquid.
- 4) Block wells by adding 100  $\mu$ l of **Blocking Buffer 2** to each well. Incubate for 10 minutes at room temperature. Decant to remove supernatant. Tap plate onto clean paper towels to remove liquid.
- 5) Just before use, mix on ice 50  $\mu$ l **ELISA ECL Substrate A** and 50  $\mu$ l **ELISA ECL Substrate B** per well of the reaction, then add 100  $\mu$ l to each well. Discard any unused chemiluminescent reagent after use.
- 6) Immediately read sample in a luminometer or microtiter-plate capable of reading chemiluminescence. "Blank" value is subtracted from all readings.

### **Reading Chemiluminescence:**

Chemiluminescence is the emission of light (luminescence) which results from a chemical reaction. The detection of chemiluminescence requires no wavelength selection because the method used is emission photometry and is not emission spectrophotometry.

To properly read chemiluminescence, make sure the plate reader is set for LUMINESCENCE mode. Typical integration time is 1 second, delay after plate movement is 100 msec. Do not use a filter when measuring light emission. Typical settings for the Synergy 2 BioTek plate reader are: use the "hole" position on the filter wheel; Optics position: Top; Read type: endpoint. Sensitivity may be adjusted based on the luminescence of a control assay without binding partner (typically we set this value as 100).

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**


Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
 San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** info@bpsbioscience.com

**Example of Assay Results:**



mCTLA4[Biotinylated]:mB7-1 binding (left) and inhibition (right), measured using the using the Mouse CTLA4[Biotinylated]:B7-1 Inhibitor Screening Assay Kit, BPS Bioscience, Catalog #79515. Luminescence was measured using a Bio-Tek fluorescent microplate reader. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at [info@bpsbioscience.com](mailto:info@bpsbioscience.com).*

**RELATED PRODUCTS:**

<u>Product Name</u>	<u>Catalog #</u>	<u>Size</u>
Mouse CTLA4 (Mouse), Fc-Fusion (Human), Avi-Tag	79062	100 µg
Mouse CTLA4 (CD152), Fc-Biotin-labeled	79001	50 µg
Human CTLA4 (CD152), Fc fusion	71149	100 µg
Human CTLA4 (CD152) Neutralizing Antibody	71212	50 µg
Mouse B7-1 (CD86), Fc fusion	79058	100 µg
Human B7-1, Fc fusion, Biotin labeled	71114	50 µg
Human B7-1, Fc fusion (Human) HiP™	71125	100 µg
Human B7-2, Fc fusion	71150	100 µg
Human B7-2 (CD86), Fc fusion, Biotin labeled	71159	50 µg
Human CD28	71113	200 µg
Human CTLA4:B7-1[Biotinylated] Inhibitor Screening Assay Kit	72009	96 rxns
Human CD28:B7-1[Biotinylated] Inhibitor Screening Assay Kit	72007	96 rxns
Human CTLA4:B7-1 TR-FRET Assay Kit	72120	384 rxns.
Anti-Human CTLA4 Neutralizing Antibody	71212	100 µg

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
 San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** info@bpsbioscience.com

### TROUBLESHOOTING GUIDE

Problem	Possible Cause	Solution
Luminescence signal of positive control reaction is weak	CTLA4 or B7-1 has lost binding capacity	Protein loses activity upon repeated freeze/thaw cycles. Use fresh CTLA4-biotin, (BPS Bioscience #79001) and fresh B7-1 (BPS Bioscience #79058). Store proteins in single-use aliquots. Increase time of protein incubation. Increase protein concentration.
	Incorrect settings on instruments	Refer to instrument instructions for settings to increase sensitivity of light detection.
	Chemiluminescent reagents mixed too soon	Chemiluminescent solution should be used within 15 minutes of mixing. Ensure both reagents are properly mixed.
	Inaccurate pipetting/technique	Run duplicates of all reactions. Use a multichannel pipettor. Use master mixes to minimize errors.
Luminescent signal is erratic or varies widely among wells	Bubbles in wells	Pipette slowly to avoid bubble formation. Tap plate lightly to disperse bubbles; be careful not to splash between wells.
	Insufficient washes	Increase number of washes. Increase wash volume. Add Tween-20 to 0.1% in washing buffer.
Background (signal to noise ratio) is high	Sample solvent is inhibiting the protein	Run negative control assay including solvent. Maintain DMSO level at <1% Increase time of protein incubation.
	Results are outside the linear range of the assay	Use different concentrations of CTLA-4-biotin (BPS Bioscience #79001) to create a standard curve.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)