



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet

CDK2 Assay Kit

Catalog #79599

DESCRIPTION: Cyclin-dependent kinases (CDKs) are key regulators of the cell cycle. CDKs are active only when bound to their regulator proteins, cyclins. CDK activity is tightly controlled for successful cell division. Since abnormal cell division represents cancer pathology, controlling CDK activity has been shown as a promising therapeutic strategy. In particular, CDK2 plays an important role in DNA replication. The *CDK2 Assay Kit* is designed to measure CDK2/CyclinA2 activity for screening and profiling applications, using Kinase-Glo[®] MAX as a detection reagent. The *CDK2 Assay Kit* comes in a convenient 96-well format, with enough purified recombinant CDK2/CyclinA2 enzyme, CDK substrate peptide, ATP and kinase assay buffer for 100 enzyme reactions.

COMPONENTS:

Catalog #	Reagent	Amount	Storage	
41101	CDK2/CyclinA2	5 µg	-80°C	Avoid multiple freeze/ thaw cycles!
79334	5x Kinase assay buffer 1	1.5 ml	-20°C	
79686	ATP (500 µM)	100 µl	-20°C	
79598	10x CDK substrate peptide 1	500 µl	-20°C	
79696	96-well plate, white	1	Room Temp.	

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Kinase-Glo MAX (Promega #V6071)
Dithiothreitol (DTT, 1 M; optional)
Microplate reader capable of reading luminescence
Adjustable micropipettor and sterile tips
30°C incubator

APPLICATIONS: Useful for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: Up to 6 months when stored as recommended.

REFERENCE: Uzma A. et. al., *Nature Review Drug Discovery* **14**:130-146 (2015)

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

- 1) Thaw **5x Kinase assay buffer 1**, **ATP** and **10x CDK substrate peptide 1**.
(Optional: If desired, add DTT to **5x Kinase assay buffer 1** to make a 10 mM concentration; e.g. add 10 µl of 1 M DTT to 1 ml **5x Kinase assay buffer 1**)
- 2) Prepare the master mixture (25 µl per well): N wells x (6 µl **5x Kinase assay buffer 1** + 1 µl **ATP (500 µM)** + 5 µl **10x CDK substrate peptide 1** + 13 µl distilled water). Add 25 µl to every well.

	Positive Control	Test Inhibitor	Blank
5x Kinase assay buffer 1	6 µl	6 µl	6 µl
ATP (500 µM)	1 µl	1 µl	1 µl
10X CDK substrate peptide 1	5 µl	5 µl	5 µl
Water	13 µl	13 µl	13 µl
Test Inhibitor	–	5 µl	–
Inhibitor Buffer (no inhibitor)	5 µl	–	5 µl
1x Kinase buffer 1	–	–	20 µl
CDK2/CyclinA2 (2.5 ng/µl)	20 µl	20 µl	–
Total	50 µl	50 µl	50 µl

- 3) Add 5 µl of Inhibitor solution of each well labeled as "Test Inhibitor". For the "Positive Control" and "Blank", add 5 µl of the same solution without inhibitor (Inhibitor buffer).
- 4) Prepare 3 ml of **1x Kinase assay buffer 1** by mixing 600 µl of **5x Kinase assay buffer 1** with 2400 µl water. 3 ml of **1x Kinase assay buffer 1** is sufficient for 100 reactions.
- 5) To the wells designated as "Blank", add 20 µl of **1x Kinase assay buffer 1**.
- 6) Thaw **CDK2/CyclinA2** enzyme on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Calculate the amount of **CDK2/CyclinA2** required for the assay and dilute enzyme to ~2.5 ng/µl with **1x Kinase assay buffer 1**. Store remaining undiluted enzyme in aliquots at -80°C.
Note: CDK2/CyclinA2 enzyme is sensitive to freeze/thaw cycles. Avoid multiple freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

- 7) Initiate reaction by adding 20 μ l of **diluted CDK2/CyclinA2** enzyme to the wells designated "Positive Control" and "Test Inhibitor Control". Incubate at 30°C for 45 minutes.
- 8) Thaw Kinase-Glo Max reagent.
- 9) After the 45-minute reaction, add 50 μ l of Kinase-Glo Max reagent to each well. Cover plate with aluminum foil and incubate the plate at room temperature for 15 minutes.
- 10) Measure luminescence using the microplate reader. "Blank" value is subtracted from all readings.

Reading Chemiluminescence:

Chemiluminescence is the emission of light (luminescence) which results from a chemical reaction. The detection of chemiluminescence requires no wavelength selection because the method used is emission photometry and is not emission spectrophotometry.

To properly read chemiluminescence, make sure the plate reader is set for LUMINESCENCE mode. Typical integration time is 1 second, delay after plate movement is 100 msec. Do not use a filter when measuring light emission. Typical settings for the Synergy 2 BioTek plate reader are: use the "hole" position on the filter wheel; Optics position: Top; Read type: endpoint. Sensitivity may be adjusted based on the luminescence of a control assay without enzyme (typically we set this value as 100).

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.


To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [**info@bpsbioscience.com**](mailto:info@bpsbioscience.com)

Please visit our website at: [**www.bpsbioscience.com**](http://www.bpsbioscience.com)

Example of Assay Results:

CDK2/Cyclin A2
 (10 μ M ATP/0.1 mg/ml CDK substrate peptide 1)



Inhibition of CDK2/CyclinA2 enzyme by Dinaciclib, measured using the CDK2 assay kit (Cat. #79599). *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com.*

RELATED PRODUCTS:

<u>Product Name</u>	<u>Catalog #</u>	<u>Size</u>
CDK1/CyclinB1, GST-tag	40454	10 μ g
CDK1/CyclinA2, GST-tag	40100	10 μ g
CDK2/CyclinA2, GST-tag	40101	10 μ g
CDK2/CyclinE1, GST-tag	40102	10 μ g
CDK2 (no tag)/CyclinA2, His-GST-tags	41101	10 μ g
CDK3/CyclinE1, GST-tag	40103	10 μ g
CDK4, FLAG-Tag	100052	20 μ g
CDK4/CyclinD3, GST-His-Tag	40104	10 μ g
CDK4(EE,T172A)/Cyclin D1, His-tag	40094	20 μ g
CDK5/p25, GST-tag	40105	10 μ g
CDK5/p35, GST-tag	40095	10 μ g
CDK6/CyclinD1, His-tag, GST-tag	40097	10 μ g
CDK6/CyclinD3, His-tags	40206	20 μ g
CDK7/CyclinH1/MNAT1, His-tags	40098	10 μ g
CDK9/CyclinK, GST-tag	40106	10 μ g
CDK9/CyclinT1, GST-tag	40307	10 μ g

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com