



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Data Sheet


Rat PDE7A Assay Kit

Catalog # 79634
Size: 96 reactions

DESCRIPTION: Phosphodiesterases (PDEs) play an important role in the dynamic regulation of cAMP and cAMP signaling. Rat PDE7A, also known as high affinity cAMP-specific phosphodiesterase 7A, has been implicated in cardiovascular function and fertility.

The Rat PDE7A Assay Kit is designed for identification of inhibitors of Rat PDE7A using fluorescence polarization. The assay is based on the binding of a fluorescent nucleotide monophosphate generated by Rat PDE7A to the binding agent.

Phosphodiesterases catalyze the hydrolysis of the phosphodiester bond in dye-labeled cyclic monophosphates. Beads selectively bind the phosphate group in the nucleotide product. This increases the size of the nucleotide relative to unreacted cyclic monophosphate. In the polarization assay, dye molecules with absorption transition vectors parallel to the linearly-polarized excitation light are selectively excited. Dyes attached to the rapidly-rotating cyclic monophosphates will obtain random orientations and emit light with low polarization. Dyes attached to the slowly-rotating nucleotide-bead complexes will not have time to reorient and therefore will emit highly polarized light.



OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

The Rat PDE7A Assay Kit comes in a convenient 96-well format, with purified Rat PDE7A enzyme, fluorescently labeled substrate (cAMP), binding agent, and PDE assay buffer for 100 enzyme reactions. The key to the *Rat PDE7A Assay Kit* is the specific binding agent. Using this kit, only two simple steps on a microtiter plate are required for Rat PDE7A reactions. First, the fluorescently labeled cAMP is incubated with a sample containing Rat PDE7A for 1 hour. Second, a binding agent is added to the reaction mix to produce a change in fluorescent polarization that can then be measured using a fluorescence reader equipped for the measurement of fluorescence polarization.

COMPONENTS:

Catalog #	Component	Amount	Storage	
60036	Rat PDE7A recombinant enzyme	>1 µg	-80°C	(Avoid freeze/thaw cycles!)
60200	FAM-Cyclic-3', 5'-AMP (20 µM)	50 µl	-80°C	
60393	PDE assay buffer	25 ml	-20°C	
60390	Binding Agent	100 µl	+4°C	
60391	Binding Agent Diluent (cAMP)	10 ml	+4°C	
79685	Black, low binding, microtiter plate	1	Room temp.	

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

Fluorescent microplate reader capable to measure fluorescence polarization.
Adjustable micropipettor and sterile tips.
1,4-Dithiothreitol (DTT), 1 M in anhydrous DMSO.

APPLICATIONS: Great for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: 6 months from date of receipt when stored as directed.

REFERENCES:

1. Morales-Garcia JA, , *et al.* (2011). Phosphodiesterase 7 Inhibition Preserves Dopaminergic Neurons in Cellular and Rodent Models of Parkinson Disease. Amédée T, ed. *PLoS ONE*. **6(2)**:e17240.
2. Giembycz, M & J. Smith, S. (2006). Phosphodiesterase 7 (PDE7) as a therapeutic target. *Drugs of The Future - DRUG FUTURE*. **31(3)**: 207.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate.

Step 1:

- 1) Dilute 20 μ M **FAM-Cyclic-3', 5'-AMP** stock 100-fold with **PDE assay buffer** to make a 200 nM solution. Make only sufficient quantity needed for the assay; store remaining 20 μ M stock solution in aliquots at -20°C.
- 2) Dilute 1M 1,4-Dithiothreitol (DTT) 1:500 into the diluted **FAM-Cyclic-3',5'-AMP**. For example, add 10 μ l DTT (1M) to 5 ml of diluted FAM-Cyclic-3', 5'-GMP (200 nM).
- 3) Add 25 μ l of **FAM-Cyclic-3',5'-AMP** (200 nM) to each well designated "Positive Control," "Test Inhibitor," and "Substrate Control."
- 4) Add 45 μ l of **PDE assay buffer** to each well designated "Blank" and add 20 μ l of **PDE assay buffer** to each well designated "Substrate Control."
- 5) Add 5 μ l of inhibitor solution to each well designated "Test Inhibitor." For the wells labeled "Positive Control," "Substrate Control," and "Blank," add 5 μ l of the same solution without inhibitor (inhibitor buffer). If inhibitor is dissolved in DMSO, make sure final DMSO concentration in the assay does not exceed 1%.
- 6) Thaw **Rat PDE7A** on ice. Upon first thaw, briefly spin tube containing enzyme to recover the full contents of the tube. Aliquot **Rat PDE7A** enzyme into single-use aliquots. Store remaining undiluted enzyme in aliquots at -80°C immediately. *Note: Rat PDE7A is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.*

	Positive Control	Test Inhibitor	Substrate Control	"Blank" Negative Control
FAM-Cyclic-3',5'-AMP (200 nM)	25 μ l	25 μ l	25 μ l	-
PDE assay buffer	-	-	20 μ l	45 μ l
Inhibitor (in PDE assay buffer)	-	5 μ l	-	-
Inhibitor Buffer (no inhibitor)	5 μ l	-	5 μ l	5 μ l
Rat PDE7A (0.02 ng/ μ l)	20 μ l	20 μ l	-	-
Total	50 μl	50 μl	50 μl	50 μl

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
 San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

- 7) Dilute **Rat PDE7A** in **PDE assay buffer** to 0.02 ng/μl (0.4 ng/reaction)*. Initiate reaction by adding 20 μl of diluted **Rat PDE7A** (0.02 ng/μl) to the wells designated "Positive Control" and "Test Inhibitor." Discard any remaining diluted enzyme after use. *Note: *Optimal enzyme concentration may vary with the specific activity of the enzyme.*
- 8) Incubate the plate at room temperature for 1 hour.

Step 2:

- 1) Mix **Binding Agent** thoroughly and dilute **Binding Agent** 1:100 with **Binding Agent Diluent**.
- 2) Add 100 μl of diluted **Binding Agent** to each microwell. Incubate at room temperature for 30 minutes with slow shaking.
- 3) Read the fluorescent polarization of the sample in a microtiter-plate reader equipped for the measurement of fluorescence polarization, capable of excitation at wavelengths ranging from 485 ± 5 nm and detection of emitted light ranging from 528 ± 10 nm. Blank value is subtracted from all other values.

CALCULATING RESULTS:

Definition of Fluorescence Polarization

$$P = \frac{I_{\parallel} - I_{\perp}}{I_{\parallel} + I_{\perp}}$$

where I_{\parallel} = Intensity with polarizers parallel and I_{\perp} = Intensity with polarizers perpendicular.

The equation above assumes that light is transmitted equally well through both parallel and perpendicular oriented polarizers. In practice, this is generally not true and a correction must be made to measure the absolute polarization state of the molecule. This correction factor is called the "G Factor".

$$FP(\text{measured}) = \frac{([I_{\parallel}] - G*[I_{\perp}])}{([I_{\parallel}] + G*[I_{\perp}])} * 1000$$

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com


Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

The G-factor is instrument-dependent and may vary slightly depending upon instrument and conditions. Please check the manual of your instrument to obtain the information about the establishment of the G-factor.

EXAMPLE OF ASSAY RESULTS:



Inhibition of Rat PDE7A by BRL-50481 measured using the *RAT PDE7A Assay Kit*, BPS Bioscience #79634. Fluorescence polarization was measured at 528 nm using a Tecan M1000 fluorescent microplate reader. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

RELATED PRODUCTS :

<u>Product Name</u>	<u>Catalog #</u>	<u>Size</u>
Rat PDE7A-HEK293 Recombinant Cell Line	60408	2 vials
Rat PDE7A	60074	10 µg
Rat PDE7B	60075	10 µg
Rat PDE4B	60049	5 µg
Rat PDE4D	60054	5 µg
Rat PDE1B	60009	10 µg
Rat PDE2A	60022	5 µg
Rat PDE10A	60102	5 µg
PDE4D2	60048	5 µg
PDE4D3	60046	5 µg
PDE4D7	60047	5 µg
PDE7A/CRE Reporter - HEK293 Cell Line	60413	2 vials
PDE7A-HEK293 Recombinant Cell Line	60407	2 vials
PDE7A-HEK293 Recombinant Cell Line	60412	2 vials
PDE4A1A	60040	10 µg
PDE4B1	60041	10 µg
PDE4B2	60042	5 µg
Dog PDE4B	60055	5 µg
Rat PDE4B Assay Kit	79571	96 rxns.
PDE Assay Kit	60300	96 rxns.
PDE4B2 Assay Kit	60343	96 rxns.
PDE4A1A Assay Kit	60340	96 rxns.
PDE4D2 Assay Kit	60345	96 rxns.
PDE4D Cell-Based Activity Assay Kit	60505	500 rxns.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com