



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
Tel: 1.858.202.1401  
Fax: 1.858.481.8694  
Email: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

## Data Sheet **PARP1 Chemiluminescent Assay Kit** Catalog # 80569

**DESCRIPTION:** The *PARP1 Chemiluminescent Activity Assay Kit* is designed to measure PARP1 activity for screening and profiling applications. PARP1 is known to catalyze the NAD-dependent addition of poly (ADP-ribose) to histones. The PARP1 assay kit comes in a convenient 384-well format, with purified PARP1 enzyme, histone mixture, activated DNA, and PARP1 assay buffer for 400 enzyme reactions. The key to the *PARP1 Chemiluminescent Activity Assay* is the biotinylated substrate. With this kit, only three simple steps are required for PARP1 reactions. First, histone proteins are coated on a 384-well plate. Next, the PARP1 biotinylated substrate is incubated with an assay buffer that contains the PARP1 enzyme. Finally, the plate is treated with streptavidin-HRP followed by addition of the ELISA ECL substrates to produce chemiluminescence that can then be measured using a chemiluminescence reader.

### COMPONENTS:

Catalog #	Reagent	Amount	Storage	<b>Avoid multiple freeze/ thaw cycles!</b>
80501	PARP1	2 x 5 µg	-80°C	
52029	5x histone mixture	2 x 1 ml	-80°C	
80601	10x Assay Mixture Containing Biotinylated Substrate	2 x 300 µl	-80°C	
80602	10x PARP assay buffer	2 x 1 ml	-20°C	
79743	Blocking buffer 3	2 x 25 ml	+4°C	
80605	Activated DNA (5X)	2 x 500 µl	-80°C	
80611	Streptavidin-HRP	2 x 100 µl	+4°C	
79670	ELISA ECL Substrate A (translucent bottle)	2 x 6 ml	Room Temp.	
	ELISA ECL Substrate B (brown bottle)	2 x 6 ml	Room Temp.	
	384-well plate	1	Room Temp.	

### MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

1x PBS buffer  
PBST buffer (1x PBS, containing 0.05% Tween-20)  
Luminometer or fluorescent microplate reader capable of reading chemiluminescence  
Adjustable micropipettor and sterile tips  
Rotating or rocker platform

**APPLICATIONS:** Great for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

**STABILITY:** Up to 1 year when stored as recommended.

**REFERENCE(S):** Brown JA, Marala RB. *J. Pharmacol. Toxicol. Methods* 2002 **47**:137-41.

**ASSAY PROTOCOL:**

*All samples and controls should be tested in duplicate.*

**Step 1: Coat 25 µl of histone solution onto a 384-well plate**

- 1) Dilute **5x histone mixture** 1:5 with PBS.
- 2) Add 25 µl of diluted histone solution to each well and incubate at 4°C overnight.
- 3) Wash the plate three times using 100 µl PBST buffer (1x PBS containing 0.05% Tween-20) per well.
- 4) Tap plate onto clean paper towel to remove liquid.
- 5) Block the wells by adding 100 µl of **Blocking buffer 3** to every well. Incubate at room temperature for 60-90 minutes.
- 6) Wash plate three times with 100 µl PBST buffer as described above.
- 7) Tap plate onto clean paper towel to remove liquid.

\*Note: coated plates can be stored at 4°C for 2-3 days before using. However it is strongly recommended that a freshly coated plate be used whenever possible.

**Step 2: Ribosylation reaction**

- 1) Prepare the master mixture: N wells x (1.25 µl **10x PARP buffer** + 1.25 µl **10X PARP Assay mixture** + 2.5 µl **Activated DNA** + 7.5 µl distilled water). Add 12.5 µl to every well.
- 2) Add 2.5 µl of Inhibitor solution of each well labeled as "Test Inhibitor". For the "Positive Control" and "Blank", add 2.5 µl of the same solution without inhibitor (Inhibitor buffer).  
*Note: The PARP1 Chemiluminescent Assay Kit is compatible with up to 1% final DMSO concentration. We recommend preparing the inhibitor in 10% DMSO aqueous solution and using 2.5 µl per PARP1 reaction.*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
Tel: 1.858.202.1401  
Fax: 1.858.481.8694  
Email: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

	Positive Control	Test Inhibitor	Blank
10x PARP buffer	1.25 µl	1.25 µl	1.25 µl
10x Assay mixture	1.25 µl	1.25 µl	1.25 µl
Activated DNA	2.5 µl	2.5 µl	2.5 µl
Water	7.5 µl	7.5 µl	7.5 µl
Test Inhibitor	-	2.5 µl	-
Inhibitor Buffer (no inhibitor)	2.5 µl	-	2.5 µl
1x PARP buffer	-	-	10 µl
PARP1 (1-2.0 ng/µl)*	10 µl	10 µl	-
Total	25 µl	25 µl	25 µl

\* If using a freshly coated plate, 10 ng/rxn of PARP1 gives good signal (S/B > 50). If using a plate that was coated and stored at 4°C for one or more days, we recommend using a higher amount of PARP1.

- 3) Prepare 1x PARP buffer by adding 1 part of **10x PARP buffer** to 9 parts distilled water (v/v)
- 4) To the wells designated as "Blank", add 10 µl of 1X PARP buffer.
- 5) Thaw **PARP1 enzyme** on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Calculate the amount of **PARP1** required for the assay and dilute enzyme to 1.0 - 2.0 ng/µl with 1x PARP buffer\*. Aliquot remaining **PARP1 enzyme** into single use aliquots. Store remaining undiluted enzyme in aliquots at -80°C. *Note: PARP1 enzyme is very sensitive to freeze/thaw cycles. Avoid multiple freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.*
- 6) Initiate reaction by adding 10 µl of diluted **PARP1 enzyme** to the wells designated "Positive Control" and "Test Inhibitor". Centrifuge the plate at 1000 rpm for 30 seconds and incubate at room temperature for 45 - 60 min.
- 7) Discard the reaction mixture after 1 hour, and wash plate three times with 100 µl PBST buffer and tap plate onto clean paper towel as described above.

### Step 3: Detection

- 1) Dilute **Streptavidin-HRP** 1:50 in **Blocking buffer 3**.
- 2) Add 25 µl of diluted Streptavidin-HRP to each well. Incubate for 30 min. at room temperature.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)


- 3) Wash three times with 100  $\mu$ l PBST buffer and tap plate onto clean paper towel as above as described above.
- 4) Just before use, mix on ice 25  $\mu$ l **ELISA ECL Substrate A** and 25  $\mu$ l **ELISA ECL Substrate B** and add 50  $\mu$ l per well.
- 5) Immediately read the plate in a luminometer or microtiter-plate reader capable of reading chemiluminescence. The "Blank" value is subtracted from all other values.

### Reading Chemiluminescence:

Chemiluminescence is the emission of light (luminescence) which results from a chemical reaction. The detection of chemiluminescence requires no wavelength selection because the method used is emission photometry and is not emission spectrophotometry.

To properly read chemiluminescence, make sure the plate reader is set for LUMINESCENCE mode. Typical integration time is 1 second, delay after plate movement is 100 msec. Do not use a filter when measuring light emission. Typical settings for the Synergy 2 BioTek plate reader are: use the "hole" position on the filter wheel; Optics position: Top; Read type: endpoint. Sensitivity may be adjusted based on the luminescence of a control assay without enzyme (typically we set this value as 100).

### Example of Assay Results:



PARP1 activity measured using the *PARP1 Chemiluminescent Activity Assay Kit*, (BPS Cat. #80569). Luminescence was measured using a Bio-Tek microplate reader. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at [info@bpsbioscience.com](mailto:info@bpsbioscience.com)*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

**RELATED PRODUCTS:**

<b><u>Product Name</u></b>	<b><u>Catalog #</u></b>	<b><u>Size</u></b>
PARP1 Assay Kit	80551	96 rxns
PARP2 Assay Kit	80552	96 rxns.
PARP3 Assay Kit	80553	96 rxns.
PARP5A (TNKS1) Assay Kit	80573	96 rxns.
PARP5B (TNKS2) Assay Kit	80579	96 rxns.
PARP6 Assay Kit	80556	32 rxns.
PARP1 Enzyme	80501	10 µg
PARP2 Enzyme	80502	10 µg
PARP3 Enzyme	80503	10 µg
PARP6 Enzyme	80506	10 µg
TNKS2 (PARP5A) Enzyme	80504	10 µg
TNKS2 (PARP5B/C) Enzyme	80505	10 µg
PARP7 Enzyme	80507	10 µg
PARP9 Enzyme	80509	10 µg
PARP11 Enzyme	80511	10 µg
PARP12 Enzyme	80512	10 µg

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)



6042 Cornerstone Court W, Ste B  
 San Diego, CA 92121  
**Tel:** 1.858.202.1401  
**Fax:** 1.858.481.8694  
**Email:** info@bpsbioscience.com

### TROUBLESHOOTING GUIDE

Problem	Possible Cause	Solution
Luminescence signal of positive control reaction is weak	PARP1 enzyme has lost activity	Enzyme loses activity upon repeated freeze/thaw cycles. Use fresh enzyme (PARP1, BPS Bioscience #80501). Store enzyme in single-use aliquots. Increase time of enzyme incubation. Increase enzyme concentration.
	Incorrect settings on instruments	Refer to instrument instructions for settings to increase sensitivity of light detection. See section on "Reading Chemiluminescence" above.
	Chemiluminescent reagents mixed too soon	Chemiluminescent solution should be used within 15 minutes of mixing. Ensure both reagents are properly mixed.
Luminescent signal is erratic or varies widely among wells	Inaccurate pipetting/technique	Run duplicates of all reactions. Use a multichannel pipettor. Use master mixes to minimize errors.
	Bubbles in wells	Pipette slowly to avoid bubble formation. Tap plate lightly to disperse bubbles; be careful not to splash between wells.
Background (signal to noise ratio) is high	Insufficient washes	Be sure to include blocking steps after wash steps. Increase number of washes. Increase wash volume. Increase Tween-20 concentration to 0.1% in TBST.
	Sample solvent is inhibiting the enzyme	Run negative control assay including solvent. Maintain DMSO level at <1% Increase time of enzyme incubation.
	Results are outside the linear range of the assay	Use different concentrations of enzyme (PARP1, BPS Bioscience #80501) to create a standard curve.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: [info@bpsbioscience.com](mailto:info@bpsbioscience.com)

Please visit our website at: [www.bpsbioscience.com](http://www.bpsbioscience.com)