



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Description**

The Chemi-Verse™ TYK2 Kinase Assay Kit is designed to measure TYK2 (Tyrosine Kinase 2) tyrosine kinase activity for screening and profiling applications using ADP-Glo™ as a detection reagent. The assay kit comes in a convenient 96-well format, with enough purified TYK2, kinase substrate, ATP, and kinase assay buffer for 100 enzyme reactions.

**Background**

Janus kinase (JAK) is a family of intracellular, nonreceptor tyrosine kinases that play an important role in cytokine-mediated signaling. This family includes four kinases: JAK1, JAK2, JAK3 and TYK2 (Tyrosine Kinase 2). Various studies have suggested that regulation of JAK activity is a promising approach to combat numerous diseases such as immune deficiency and cancer. Upon binding of a cytokine to its corresponding receptor, the receptor changes conformation and recruits two JAK molecules, which phosphorylate the receptor and activate a signal that results in STAT dimerization and translocation to the nucleus. This results in expression of pro-inflammatory cytokines and growth factors. For example, IL-12 and IL-23 receptor pathways require a TYK2/JAK2 heterodimer. TYK2 is also required for type interferon and IL-10 signaling. TYK2 has been closely linked to psoriasis, with individuals that have loss-of-function mutations being protected from this disorder. This points to the possibility of inhibiting TYK2 as a treatment option. Several TYK2 inhibitors have been developed for the treatment of psoriasis, hidradenitis suppurative, IBD (inflammatory bowel disease), dermatomyositis and SLE (systemic lupus erythematosus). Deucravacitinib, an allosteric inhibitor that binds to the pseudokinase domain, has been approved in 2022 for the treatment of plaque psoriasis. This approval indicates the potential of targeting TYK2 to treat inflammatory disorders.

**Applications**

Study enzyme kinetics and screen small molecule inhibitors for drug discovery and high throughput screening (HTS) applications.

**Supplied Materials**

Catalog #	Name	Amount	Storage
40285	Tyk2, His-Tag*	2 µg	-80°C
79334	5x Kinase Buffer 1	1.5 ml	-20°C
79686	500 µM ATP	50 µl	-20°C
79519	10x IRS1-tide	500 µl	-20°C
82545	White 96-well plate	1	Room Temperature

\*The concentration of the protein is lot-specific and will be indicated on the tube.

**Materials Required but Not Supplied**

Name	Ordering Information
ADP-Glo™ Kinase Assay	Promega #V6930
DTT (Dithiothreitol), 1M, optional	
Microplate reader capable of reading luminescence	
Adjustable micropipettor and sterile tips	
30°C incubator	

**Storage Conditions**

This assay kit will perform optimally for up to **6 months** from date of receipt when the materials are stored as directed.

**Safety**

This product is for research purposes only and not for human or therapeutic use. This product should be considered hazardous and is harmful by inhalation, in contact with skin, eyes, clothing, and if swallowed. If contact occurs, wash thoroughly.

**Assay Principle**

The **ADP-Glo™ Kinase Assay (Promega #V6930)** quantifies the amount of ADP produced by a kinase upon phosphorylation of a substrate. First, addition of the ADP-Glo™ reagent terminates the reaction and quenches the remaining ATP. Second, the addition of the Kinase Detection reagent converts the produced ADP to ATP. The newly generated ATP is quantified by a luciferase reaction. The luminescent signal correlates with the amount of ADP generated by the kinase and is linear to 1 mM ATP.

**Contraindications**

The final concentration of DMSO in the assay should not exceed 1%.

**Assay Protocol**

- All samples and controls should be tested in duplicate.
- The assay should include “Blank”, “Positive Control” and “Test Inhibitor” conditions.
- We recommend maintaining the diluted protein on ice during use.
- For detailed information on protein handling please refer to [Protein FAQs \(bpsbioscience.com\)](https://www.bpsbioscience.com).
- We recommend using Ilginatinib (#82587) or Staurosporine (#27002) as internal control. If not running a dose response curve for the control inhibitor, we recommend running the control inhibitor at 0.1X, 1X and 10X the IC<sub>50</sub> value shown in the validation data below.
- For instructions on how to prepare reagent dilutions please refer to [Serial Dilution Protocol \(bpsbioscience.com\)](https://www.bpsbioscience.com).

1. Thaw **5x Kinase Assay Buffer 1**, **500 μM ATP**, and **10x IRS1-tide substrate (1 mg/ml)**.

*Optional: If desired, make 5x Kinase Assay Buffer 1 with 10 mM DTT.*

2. Prepare 3 ml of **1x Kinase Assay Buffer 1** by mixing 600 μl of **5x Kinase Assay Buffer 1** with 2,400 μl of distilled water.

*Note: Three (3 ml) of 1x Kinase Assay Buffer 1 is sufficient for 100 reactions.*

3. Prepare a **Master Mix** (12.5 μl/well): N wells x (6 μl of 5x Kinase Assay Buffer 1 + 0.5 μl of 500 μM ATP + 2.5 μl of 10x IRS1-tide substrate (1 mg/ml) + 3.5 μl of distilled water).
4. Add 12.5 μl of Master Mix to every well.

- Prepare the **Test Inhibitor** (2.5 µl/well): for a titration prepare serial dilutions at concentrations 10-fold higher than the desired final concentrations. The final volume of the reaction is 25 µl.

5.1 If the Test Inhibitor is water-soluble: Prepare serial dilutions in 1x Kinase Assay Buffer 1, 10-fold more concentrated than the desired final concentrations.

For the positive and negative controls, use 1x Kinase Assay Buffer 1 (Diluent Solution).

**OR**

5.2 If the Test inhibitor is soluble in DMSO: Prepare the test inhibitor at 100-fold the highest desired concentration in 100% DMSO, then dilute the inhibitor 10-fold in 1x Kinase Assay Buffer 1 to prepare the highest concentration of the 10-fold intermediate dilutions. The concentration of DMSO is now 10%.

Prepare serial dilutions of the Test Inhibitor at 10-fold the desired final concentrations using 10% DMSO in 1x Kinase Assay Buffer 1 to keep the concentration of DMSO constant.

For positive and negative controls, prepare 10% DMSO in 1x Kinase Assay Buffer 1 (vol/vol) so that all wells contain the same amount of DMSO (Diluent Solution).

*Note: The final concentration of DMSO should not exceed 1%.*

- Add 2.5 µl of Test Inhibitor to each well labeled "Test Inhibitor".
- Add 2.5 µl of Diluent Solution to the "Positive Control" and "Blank" wells.
- Add 10 µl of 1x Kinase Assay Buffer 1 to the "Blank" wells.
- Thaw **TYK2 Kinase** on ice. Briefly spin the tube to recover its full content.
- Dilute the protein kinase (10 µl/well) to 2 ng/µl with **1x Kinase Assay Buffer 1**.
- Initiate the reaction by adding 10 µl of diluted kinase to the wells designated "Positive Control" and "Test Inhibitor".

Component	Blank	Positive Control	Test Inhibitor
Master Mix	12.5 µl	12.5 µl	12.5 µl
Test Inhibitor	-	-	2.5 µl
Diluent Solution	2.5 µl	2.5 µl	-
1x Kinase Assay Buffer 1	10 µl	-	-
Diluted TYK2 (2 ng/µl)	-	10 µl	10 µl
<b>Total</b>	<b>25 µl</b>	<b>25 µl</b>	<b>25 µl</b>

- Incubate at 30°C for 45 minutes.


13. Thaw the ADP-Glo™ reagent.
14. At the end of the 45 minute reaction, add 25 µl of ADP-Glo™ reagent to each well.
15. Cover the plate with aluminum foil and incubate at Room Temperature (RT) for 45 minutes.
16. Thaw the Kinase Detection Reagent.
17. Add 50 µl of Kinase Detection reagent to each well.
18. Cover the plate with aluminum foil and incubate at RT for another 45 minutes.
19. Immediately read in a luminometer or a microplate reader capable of reading luminescence.
20. The “Blank” value is subtracted from all other readings.

### Reading Luminescence

Luminescence is the emission of light resulting from a chemical reaction. The detection of luminescence requires no wavelength selection because the method used is emission photometry and not emission spectrophotometry.

To properly read luminescence, make sure the plate reader is set for LUMINESCENCE mode. Typical integration time is 1 second, delay after plate movement is 100 msec. Do not use a filter when measuring light emission. Typical settings for the Synergy 2 BioTek plate reader: use the “hole” position on the filter wheel; Optics position: Top; Read type: endpoint. Sensitivity may be adjusted based on the luminescence of a control assay without enzyme (typically we set this value as 100).

## Example Results



*Figure 1: Inhibition of TYK2 kinase activity by the inhibitors Ilginatinib or Staurosporine.* TYK2 kinase activity was measured in the presence of increasing concentrations of Ilginatinib (#82587) or Staurosporine (#27002). The “Blank” value was subtracted from all other values. Results are expressed as the percent of control (kinase activity in the absence of inhibitor, set at 100%).

*Data shown is representative. For lot-specific information, please contact BPS Bioscience, Inc. at [support@bpsbioscience.com](mailto:support@bpsbioscience.com).*

## Troubleshooting Guide

Visit [bpsbioscience.com/assay-kits-faq](https://bpsbioscience.com/assay-kits-faq) for detailed troubleshooting instructions. For all further questions, please email [support@bpsbioscience.com](mailto:support@bpsbioscience.com)

## References

Rusinol L. and Puig L., 2023 *Int J Mol Sci* 24(4) :3391.  
Morand E., *et al.*, 2024 *Nature Reviews Rheumatology* 20 :232-240.

## Related Products

Products	Catalog #	Size
TYK2 (JH2 Pseudokinase Domain) Inhibitor Screening Assay Kit	78107	384 reactions
Tyk2, GST-Tag Recombinant	100400	10 µg
TYK2 (JH2 Domain), FLAG-Tag Recombinant	100523	10 µg
Chemi-Verse™ JAK1 Kinase Assay Kit	82542	96 reactions
Chemi-Verse™ JAK3 Kinase Assay Kit	82544	96 reactions

Version 062024